

# Conformal foliations

Michael Eastwood

[joint work with Paul Baird]


Australian National University

# Disclaimers and references

- R.P. Kerr, I. Robinson, . . . ~1961
- R. Penrose and W. Rindler, *Spinors and Space-time* vol. 2, Chapter 7, Cambridge University Press 1986
- P. Baird and J.C. Wood, *Harmonic morphisms and shear-free ray congruences* (2002),  
<http://www.maths.leeds.ac.uk/pure/staff/wood/BWBook/BWBook.html>
- P. Baird and J.C. Wood, *Harmonic Morphisms between Riemannian Manifolds*, Oxford University Press 2003
- P. Baird and R. Pantilie, *Harmonic morphisms on heaven spaces*, *Bull. Lond. Math. Soc.* **41** (2009) 198–204
- P. Nurowski, *Construction of conjugate functions*, *Ann. Glob. Anal. Geom.* **37** (2010) 321–326
- P. Baird and M.G. Eastwood, *CR geometry and conformal foliations*,  
<http://arxiv.org/abs/1011.4717>

# Conformal foliations

$U$  = unit vector field on  $\Omega^{\text{open}} \subseteq \mathbb{R}^3$ .



$U$  is (transversally) conformal  
 $\Leftrightarrow \mathcal{L}_U$  preserves the conformal metric orthogonal to its leaves

$h$   
 $\mathbb{C}$

isothermal  
coördinates



$$h = f + ig \quad \langle \nabla f, \nabla g \rangle = 0$$
$$\|\nabla f\| = \|\nabla g\|$$

conjugate functions

# Conjugate functions on $\mathbb{R}^2$

$$f = f(r, s) \quad g = g(r, s) \quad \text{s.t.} \quad \begin{cases} \langle \nabla f, \nabla g \rangle = 0 \\ \|\nabla f\| = \|\nabla g\| \end{cases}$$

- $f = r \quad g = s$
- $f = r^2 - s^2 \quad g = 2rs$
- $f = \frac{r}{r^2 + s^2} \quad g = \frac{s}{r^2 + s^2}$
- $f = e^r \cos s \quad g = e^r \sin s$

$h \equiv f + ig$  is (anti-)holomorphic in  $z \equiv r + is$

# Conjugate functions on $\mathbb{R}^3$

$$f = f(q, r, s) \quad g = g(q, r, s) \quad \text{s.t.} \quad \begin{cases} \langle \nabla f, \nabla g \rangle = 0 \\ \|\nabla f\| = \|\nabla g\| \end{cases}$$

- $f = r \quad g = s$

- $f = q^2 - r^2 - s^2 \quad g = 2q\sqrt{r^2 + s^2}$

- $f = r \frac{q^2 + r^2 + s^2}{r^2 + s^2} \quad g = s \frac{q^2 + r^2 + s^2}{r^2 + s^2}$

- $f = \frac{(1 - q^2 - r^2 - s^2)r + 2qs}{r^2 + s^2}$   
 $g = \frac{(1 - q^2 - r^2 - s^2)s - 2qr}{r^2 + s^2}$

$$\begin{array}{ccc} \mathbb{R}^3 & \hookrightarrow & S^3 \\ & & \downarrow \text{Hopf} \\ \mathbb{R}^2 & \leftarrow & S^2 \setminus \{*\} \end{array}$$

# Almost Hermitian structures

NB:  $J(p, q, r, s) : \mathbb{R}^4 \rightarrow \mathbb{R}^4$  satisfies

- $J^2 = -\text{Id}$
  - $J \in \text{SO}(4)$
- $$\iff J = \begin{bmatrix} 0 & -u & -v & -w \\ u & 0 & -w & v \\ v & w & 0 & -u \\ w & -v & u & 0 \end{bmatrix}$$
- $u^2 + v^2 + w^2 = 1$ , two-sphere

Consider  $\mathbb{R}^3 = \{(p, q, r, s) \in \mathbb{R}^4 \mid p = 0\} \subset \mathbb{R}^4$

NB:  $U \equiv \left( J \frac{\partial}{\partial p} \right) \Big|_{\mathbb{R}^3} = \left( u \frac{\partial}{\partial q} + v \frac{\partial}{\partial r} + w \frac{\partial}{\partial s} \right) \Big|_{\mathbb{R}^3}$

unit vector field

also  $\leadsto$  two-sphere

# Sphere bundles

bundle of  
unit vectors

bundle of almost  
Hermitian structures

$$\begin{array}{ccc}
 Q_0 & \subset & Z_0 \\
 \downarrow & & \downarrow \\
 \mathbb{R}^3 & \subset & \mathbb{R}^4
 \end{array}$$

$\tau$

section



unit vector field

section



almost Hermitian structure

# Hermitian structures

Lemma

$J$  is integrable  $\implies U \equiv \left( J \frac{\partial}{\partial p} \right) \Big|_{\mathbb{R}^3}$  is conformal

Conversely??

NB:  $J$  integrable  $\implies J$  real-analytic

Question:  $U$  conformal  $\implies U$  real-analytic??

Answer: NO!

However:  $U$  real-analytic and conformal  
 $\implies U$  extends uniquely to an integrable  $J$ .

} WHY?



# Twistor geometry

bundle of almost  
Hermitian structures

$$\begin{array}{ccc}
 Q_0 \subset Z_0 = \mathbb{C}P_3 \setminus \{z_3 = z_4 = 0\} \ni [z_1, z_2, z_3, z_4] & & \\
 \downarrow & \tau \downarrow & \downarrow \\
 \mathbb{R}^3 \subset \mathbb{R}^4 = \mathbb{C}^2 \ni \begin{pmatrix} p + iq \\ r + is \end{pmatrix} = \frac{1}{|z_3|^2 + |z_4|^2} \begin{pmatrix} z_2 \bar{z}_3 + z_4 \bar{z}_1 \\ z_1 \bar{z}_3 - z_4 \bar{z}_2 \end{pmatrix} & & 
 \end{array}$$

compactify


$\mathbb{C}P_3$

$\tau \downarrow$


$S^4$

twistor fibration (cf. Hopf)

# Twistor geometry cont'd




$$\begin{aligned}
 Q &= \{ [z] \in \mathbb{C}P_3 \mid \Re(z_2 \bar{z}_3 + z_4 \bar{z}_1) = 0 \} \\
 &\cong \{ [Z] \in \mathbb{C}P_3 \mid |Z_1|^2 + |Z_2|^2 = |Z_3|^2 + |Z_4|^2 \} \\
 &\equiv \underline{\text{Levi-indefinite hyperquadric}}
 \end{aligned}$$



(cf. saddle)

# Twistor results

$$\begin{array}{ccc}
 Q & \subset & \mathbb{C}P_3 \\
 \downarrow & & \downarrow \tau \\
 S^3 & \subset & S^4
 \end{array}$$



Theorem A section  $S^4 \supseteq \text{open } \Omega \xrightarrow{J} \mathbb{C}P_3$  of  $\tau$  defines an integrable Hermitian structure if and only if  $\tilde{M} \equiv J(\Omega)$  is a complex submanifold.

Theorem A section  $S^3 \supseteq \text{open } \Omega \xrightarrow{U} Q$  of  $\tau : Q \rightarrow S^3$  defines a conformal foliation if and only if  $M \equiv U(\Omega)$  is a CR submanifold.

# CR submanifolds and functions

$M \subset Q \subset \mathbb{C}\mathbb{P}_3$  is a ‘CR submanifold’?

It means:  $TM \cap JTQ$  is preserved by  $J$ .

It does not mean:  $M = \{f = 0\}$  where  $f$  is a


CR function:  $(X + iJX)f = 0 \quad \forall X \in \Gamma(TQ \cap JTQ)$ .

Implicit function theorem  
is false in the CR category

- CR functions on  $Q$  are real-analytic.
- conformal foliations on  $S^3$  need not be.

# CR functions


$$\{[Z] \in \mathbb{C}\mathbb{P}_2 \mid |Z_1|^2 + |Z_2|^2 = |Z_3|^2\} = \text{three-sphere}$$



Theorem (H. Lewy 1956)

CR  $\Rightarrow$  holomorphic extension

$$\{[Z] \in \mathbb{C}\mathbb{P}_3 \mid |Z_1|^2 + |Z_2|^2 = |Z_3|^2 + |Z_4|^2\} = Q$$



Corollary


CR  $\Rightarrow$  holomorphic extension

Hence, a CR function on  $Q$  is real-analytic!

# Smooth conjugate functions

Eikonal equation:  $\left(\frac{\partial f}{\partial r}\right)^2 + \left(\frac{\partial f}{\partial s}\right)^2 = 1$

Plenty of non-analytic solutions:



$f =$  signed distance to  $\Gamma$

$$\left. \begin{array}{l} f(q, r, s) = f(r, s) \\ g(q, r, s) = q \end{array} \right\} \Rightarrow \left\{ \begin{array}{l} \langle \nabla f, \nabla g \rangle = 0 \\ \|\nabla f\| = \|\nabla g\| \end{array} \right.$$

QED

THANK YOU