

# Prime Numbers

## How Far Apart Are They?

Stijn S.C. Hanson

February 19, 2015

## 1 Distribution of Prime Numbers

- Behaviour of  $\pi(x)$
- Behaviour of  $\pi(x; a, q)$

## 2 Prime Constellations

- Distance Between Neighbouring Primes
- Beyond Bounded Gaps

## 1 Distribution of Prime Numbers

- Behaviour of  $\pi(x)$
- Behaviour of  $\pi(x; a, q)$

## 2 Prime Constellations

- Distance Between Neighbouring Primes
- Beyond Bounded Gaps

# Basic Definitions

- A prime number is any positive integer bigger than 1 whose only divisors are 1 and itself;

# Basic Definitions

- A prime number is any positive integer bigger than 1 whose only divisors are 1 and itself;
- We define  $\pi(x)$  to be the number of prime numbers less than or equal to  $x$  and  $p_n$  to be the  $n$ th prime number;

# Basic Definitions

- A prime number is any positive integer bigger than 1 whose only divisors are 1 and itself;
- We define  $\pi(x)$  to be the number of prime numbers less than or equal to  $x$  and  $p_n$  to be the  $n$ th prime number;
- Two integers  $n$  and  $m$  are said to be coprime if their greatest common divisor is 1. I.e. if  $(n, m) = 1$ ;

# Basic Definitions

- A prime number is any positive integer bigger than 1 whose only divisors are 1 and itself;
- We define  $\pi(x)$  to be the number of prime numbers less than or equal to  $x$  and  $p_n$  to be the  $n$ th prime number;
- Two integers  $n$  and  $m$  are said to be coprime if their greatest common divisor is 1. I.e. if  $(n, m) = 1$ ;
- The number of coprime integers to  $n$  which do not exceed  $n$  is denoted by  $\varphi(n)$ .

# Growth of $\pi(x)$

Theorem (Prime Number Theorem, Hadamard-de la Vallée Poussin, 1896)

$$\pi(x) = \frac{x}{\log x} + O\left(\frac{x}{(\log x)^2}\right)$$



# Growth of $\pi(x)$

Theorem (Prime Number Theorem, Hadamard-de la Vallée Poussin, 1896)

$$\pi(x) = \frac{x}{\log x} + O\left(\frac{x}{(\log x)^2}\right)$$

$$\pi(x) = \int_2^{\infty} \frac{dx}{\log x} + O\left(\frac{x}{\log x} e^{-a\sqrt{\log x}}\right)$$

Growth of  $\pi(x)$ 

Theorem (Prime Number Theorem, Hadamard-de la Vallée Poussin, 1896)

$$\pi(x) = \frac{x}{\log x} + O\left(\frac{x}{(\log x)^2}\right)$$

$$\pi(x) = \int_2^{\infty} \frac{dx}{\log x} + O\left(\frac{x}{\log x} e^{-a\sqrt{\log x}}\right)$$

$$\pi(x) = \int_2^{\infty} \frac{dx}{\log x} + O(\sqrt{x} \log x) \quad (RH)$$

Growth of  $\pi(x)$ 

Theorem (Prime Number Theorem, Hadamard-de la Vallée Poussin, 1896)

$$\pi(x) = \frac{x}{\log x} + O\left(\frac{x}{(\log x)^2}\right)$$

$$\pi(x) = \int_2^{\infty} \frac{dx}{\log x} + O\left(\frac{x}{\log x} e^{-a\sqrt{\log x}}\right)$$

$$\pi(x) = \int_2^{\infty} \frac{dx}{\log x} + O(\sqrt{x} \log x) \quad (RH)$$

Putting  $x + \log x$  into the top characterisation tells us that the asymptotic gap between primes is  $\log x$ .

# Primes in Short Intervals

## Theorem (Heath-Brown, 1988)

Let  $\theta \in (7/12, 1)$ . Then

$$\pi(x + x^\theta) - \pi(x) \sim \frac{x^\theta}{\log x}$$

# Primes in Short Intervals

## Theorem (Heath-Brown, 1988)

Let  $\theta \in (7/12, 1)$ . Then

$$\pi(x + x^\theta) - \pi(x) \sim \frac{x^\theta}{\log x}$$

## Theorem (Baker-Harman-Pintz, 2001)

Every interval of the form  $[x - x^{0.525}, x]$  contains prime numbers if  $x$  is big enough..

# Arithmetic Progressions

## Theorem (Dirichlet, 1837)

*Let  $a, q \in \mathbb{N}$  be two coprime integers. Then the arithmetic progression*

$$a, a + q, a + 2q, \dots$$

*contains infinitely many prime numbers.*

# Arithmetic Progressions

## Theorem (Dirichlet, 1837)

*Let  $a, q \in \mathbb{N}$  be two coprime integers. Then the arithmetic progression*

$$a, a + q, a + 2q, \dots$$

*contains infinitely many prime numbers.*

## Theorem (Green-Tao, 2004)

*The primes contain arbitrarily long arithmetic progressions.*

Growth of  $\pi(x; a, q)$ 

## Theorem (Barban-Bombieri-Vinogradov, 1961-1987-1965)

For any small  $\epsilon$  and real  $A$

$$\sum_{q \leq x^{1/2-\epsilon}} \sup_{a \in (\mathbb{Z}/q\mathbb{Z})^*} \left| \pi(x; a, q) - \frac{1}{\varphi(q)} \pi(x) \right| \ll x(\log x)^{-A}.$$



# Growth of $\pi(x; a, q)$

## Theorem (Barban-Bombieri-Vinogradov, 1961-1987-1965)

For any small  $\epsilon$  and real  $A$

$$\sum_{q \leq x^{1/2-\epsilon}} \sup_{a \in (\mathbb{Z}/q\mathbb{Z})^*} \left| \pi(x; a, q) - \frac{1}{\varphi(q)} \pi(x) \right| \ll x(\log x)^{-A}.$$

## Conjecture (Elliott-Halberstam, 1968)

For all  $\theta \in (0, 1)$

$$\sum_{q \leq x^\theta} \sup_{a \in (\mathbb{Z}/q\mathbb{Z})^*} \left| \pi(x; a, q) - \frac{1}{\varphi(q)} \pi(x) \right| \ll x(\log x)^{-A}.$$

If this holds for some  $\theta \in (0, 1)$  then we say that the primes have level of distribution  $\theta$ .

## 1 Distribution of Prime Numbers

- Behaviour of  $\pi(x)$
- Behaviour of  $\pi(x; a, q)$

## 2 Prime Constellations

- Distance Between Neighbouring Primes
- Beyond Bounded Gaps

# Normalised Prime Gaps

Theorem (Westzynthius, 1931)

Let  $p_n$  be the  $n$ th prime number. Then

$$\limsup_{n \rightarrow \infty} \frac{p_{n+1} - p_n}{\log n} = \infty.$$

# Normalised Prime Gaps

## Theorem (Westzynthius, 1931)

Let  $p_n$  be the  $n$ th prime number. Then

$$\limsup_{n \rightarrow \infty} \frac{p_{n+1} - p_n}{\log n} = \infty.$$

## Theorem (Golston-Pintz-Yildirim, 2006)

$$\liminf_{n \rightarrow \infty} \frac{p_{n+1} - p_n}{\log n} = 0$$

# Normalised Prime Gaps

## Theorem (Westzynthius, 1931)

Let  $p_n$  be the  $n$ th prime number. Then

$$\limsup_{n \rightarrow \infty} \frac{p_{n+1} - p_n}{\log n} = \infty.$$

## Theorem (Golston-Pintz-Yildirim, 2006)

$$\liminf_{n \rightarrow \infty} \frac{p_{n+1} - p_n}{\log n} = 0$$

## Theorem (Banks-Freiman-Maynard, 2014)

The set of limit points of the sequence of normalised prime gaps contains 2% of all non-negative real numbers.

# Bounded Prime Gaps

## Conjecture (Twin Prime Conjecture)

*There are infinitely many prime numbers  $p$  and  $q$  which differ by precisely 2. In other words*

$$\liminf_{n \rightarrow \infty} (p_{n+1} - p_n) = 2$$

# Bounded Prime Gaps

## Conjecture (Twin Prime Conjecture)

*There are infinitely many prime numbers  $p$  and  $q$  which differ by precisely 2. In other words*

$$\liminf_{n \rightarrow \infty} (p_{n+1} - p_n) = 2$$

## Theorem (Zhang, 2013)

$$\liminf_{n \rightarrow \infty} (p_{n+1} - p_n) \leq 70,000,000$$

# Bounded Prime Gaps

## Conjecture (Twin Prime Conjecture)

*There are infinitely many prime numbers  $p$  and  $q$  which differ by precisely 2. In other words*

$$\liminf_{n \rightarrow \infty} (p_{n+1} - p_n) = 2$$

## Theorem (Maynard-Zhang, 2013)

$$\liminf_{n \rightarrow \infty} (p_{n+1} - p_n) \leq 600$$



# Bounded Prime Gaps

## Conjecture (Twin Prime Conjecture)

*There are infinitely many prime numbers  $p$  and  $q$  which differ by precisely 2. In other words*

$$\liminf_{n \rightarrow \infty} (p_{n+1} - p_n) = 2$$

## Theorem (Maynard-Polymath-Zhang, 2013-2014)

$$\liminf_{n \rightarrow \infty} (p_{n+1} - p_n) \leq 246$$

# Bounded Prime Gaps

## Conjecture (Twin Prime Conjecture)

*There are infinitely many prime numbers  $p$  and  $q$  which differ by precisely 2. In other words*

$$\liminf_{n \rightarrow \infty} (p_{n+1} - p_n) = 2$$

## Theorem (Maynard-Polymath-Zhang, 2013-2014)

$$\liminf_{n \rightarrow \infty} (p_{n+1} - p_n) \leq 246$$

## Theorem (Maynard, 2014)

$$\liminf_{n \rightarrow \infty} (p_{n+m} - p_n) \ll m^3 e^{4m}$$

# Generalisations of the Bounded Gaps Conjecture

## Conjecture (Polignac, 1843)

*Let  $k$  be any positive integer. Then, for infinitely many  $n \in \mathbb{N}$ , we have that  $p_{n+1} - p_n = 2k$ . If this holds then  $2k$  is called a Polignac number.*

# Generalisations of the Bounded Gaps Conjecture

## Conjecture (Polignac, 1843)

*Let  $k$  be any positive integer. Then, for infinitely many  $n \in \mathbb{N}$ , we have that  $p_{n+1} - p_n = 2k$ . If this holds then  $2k$  is called a Polignac number.*

## Conjecture (Dickson, 1904)

*Let  $a_1 + b_1n, a_2 + b_2n, \dots, a_k + b_kn$  be a finite set of linear forms with integer coefficients where  $b_i \geq 1$  for all  $1 \leq i \leq k$ . Then, if there is no positive integer  $m$  divide all the products  $\prod_{i=1}^k f_i(n)$  for all integers  $n$  then there exist infinitely many natural numbers  $n$  such that all of the linear forms are prime.*

# Partial Results Towards Polignac's Conjecture

## Theorem (Pintz, 2013)

*There is some ineffective constant  $c$  such that every interval of the form  $[m, m + c]$  contains a Polignac number.*

# Partial Results Towards Polignac's Conjecture

## Theorem (Pintz, 2013)

*There is some ineffective constant  $c$  such that every interval of the form  $[m, m + c]$  contains a Polignac number.*

## Corollary

*The set of Polignac numbers contains arbitrarily long arithmetic progressions.*

# Partial Results Towards Polignac's Conjecture

## Theorem (Pintz, 2013)

*There is some ineffective constant  $c$  such that every interval of the form  $[m, m + c]$  contains a Polignac number.*

## Corollary

*The set of Polignac numbers contains arbitrarily long arithmetic progressions.*

## Theorem (Hanson, 2014)

*Every arithmetic progression of the form  $q, 2q, 3q, \dots$  contains infinitely many Polignac numbers.*