

# DIFFERENTIAL CHARACTERS OF DRINFELD MODULES AND DE RHAM COHOMOLOGY

JAMES BORGER AND ARNAB SAHA

ABSTRACT. We introduce differential characters of Drinfeld modules. These are function-field analogues of Buium's  $p$ -adic differential characters of elliptic curves and of Manin's differential characters of elliptic curves in differential algebra, both of which have had notable Diophantine applications. We determine the structure of the group of differential characters. This shows the existence of a family of interesting differential modular functions on the moduli of Drinfeld modules. It also leads to a canonical  $F$ -crystal equipped with a map to the de Rham cohomology of the Drinfeld module. This  $F$ -crystal is of a differential-algebraic nature, and the relation to the classical cohomological realizations is presently not clear.

## 1. INTRODUCTION

The theory of arithmetic jet spaces developed by Buium draws inspiration from the theory of differential algebra over a function field. In differential algebra, given a scheme  $E$  defined over a function field  $K$  with a derivation  $\partial$  on it, one can define the jet spaces  $J^n E$  for all  $n \in \mathbb{N}$  with respect to  $(K, \partial)$  and they form an inverse system of schemes satisfying a universal property with respect to derivations lifting  $\partial$ . The ring of global functions  $\mathcal{O}(J^n E)$  can be thought of as the ring of  $n$ -th order differential functions on  $E$ . In the case when  $E$  is an elliptic curve and its structure sheaf  $\mathcal{O}_E$  does not have a derivation lifting  $\partial$  (if it does, then it is the isotrivial case and  $E$  will descend to the subfield  $K^{\partial=0}$  of constants), there exists a differential function  $\Theta \in \mathcal{O}(J^2 E)$  which is a homomorphism of group schemes from  $J^2 E$  to the additive group  $\mathbb{G}_a$ . Such a  $\Theta$  is an example of a differential character of order 2 for  $E$  and is known as a Manin character. Explicitly, if  $E$  is given by the Legendre equation  $y^2 = x(x-1)(x-t)$  over  $K = \mathbb{C}(t)$  with derivation  $\partial = \frac{d}{dt}$ , then

$$\Theta(x, y, x', y', x'', y'') = \frac{y}{2(x-t)^2} - \frac{d}{dt} \left[ 2t(t-1) \frac{x'}{y} \right] + 2t(t-1) x' \frac{y'}{y^2}.$$

The existence of such a  $\Theta$  is a consequence of the Picard–Fuchs equation. Using the derivation  $\partial$  on  $K$ , we can lift any  $K$ -rational point  $P \in E(K)$  canonically to  $J^2 E(K)$ , and this defines a homomorphism  $\nabla : E(K) \rightarrow J^2 E(K)$ . We emphasize that  $\nabla$  is merely a map on  $K$ -rational points and does not come from a map of schemes. The composition  $\Theta \circ \nabla : E(K) \rightarrow \mathbb{G}_a(K)$  is then a group homomorphism of  $K$ -points. Note that the torsion points of  $E(K)$  are contained in the kernel of  $\Theta$  since  $\mathbb{G}_a(K)$  is torsion free. Such a  $\Theta$  was used by Manin to give a proof of the Lang–Mordell conjecture for abelian varieties over function fields [M]. Later Buium gave a different proof, using other methods, but still using the Manin map [Bui1].

---

*Date:* August 28, 2017.

The theory of arithmetic jet spaces, as developed by Buium, proceeds similarly. Derivations  $\partial$  are replaced by what are known as  $\pi$ -derivations  $\delta$ . They naturally arise from the theory of  $\pi$ -typical Witt vectors. For instance, when our base ring  $R$  is an unramified extension of the ring of  $p$ -adic integers  $\mathbb{Z}_p$ , for a fixed prime  $\pi = p$ , the Fermat quotient operator  $\delta x = \frac{\phi(x) - x^p}{p}$  is the unique  $p$ -derivation, where the endomorphism  $\phi: R \rightarrow R$  is the lift of the  $p$ -th power Frobenius endomorphism of  $R/pR$ . In analogy with differential algebra, one can define the  $n$ -th order jet space  $J^n E$  of an elliptic curve  $E$  over  $R$  to be the ( $\pi$ -adic) formal scheme over  $R$  with functor of points

$$(J^n E)(C) = \text{Hom}_R(\text{Spec } W_n(C), E),$$

where  $W_n(C)$  is the ring of  $\pi$ -typical Witt vectors of length  $n + 1$ , which we view as the arithmetic analogue of  $C[t]/(t^{n+1})$ . The jet space  $J^n E$  is also known as the Greenberg transform. As with the differential jet space, it has relative dimension  $n + 1$  over the base, in this case  $\text{Spf } R$ .

Then one can define  $\mathbf{X}_n(E)$  to be the  $R$ -module of all group-scheme homomorphisms from  $J^n E$  to the  $\pi$ -adic formal scheme  $\hat{\mathbb{G}}_a$ . Let  $\mathbf{X}_\infty(E)$  be the direct limit of the  $\mathbf{X}_n(E)$ . Now the usual Frobenius operator on Witt vectors induces a canonical Frobenius morphism  $\phi: J^{n+1} E \rightarrow J^n E$  lying over the endomorphism  $\phi$  of  $\text{Spf } R$ . Hence pulling back morphisms via  $\phi$  as  $\Theta \mapsto \phi^* \Theta$ , endows  $\mathbf{X}_\infty(E)$  with an action of  $\phi^*$  and hence makes  $\mathbf{X}_\infty(E)$  into a left module over the twisted polynomial ring  $R\{\phi^*\}$  with commutation law  $\phi^* \cdot r = \phi(r) \cdot \phi^*$ . In [Bui2], Buium studied the structure of  $\mathbf{X}_\infty(E)$ . Putting  $K = R[\frac{1}{p}]$ , he showed that  $\mathbf{X}_\infty(E) \otimes_R K$  is freely generated by a single element as a  $K\{\phi^*\}$ -module. This element is of order 2 unless  $E$  is a canonical lift, in which case it is of order 1. It is the arithmetic analogue of the Manin character.

In this paper, we study the function-field analogue of Buium's theory. We emphasize that we take the function-field analogue in every possible sense. So instead of looking at characters  $J^n E \rightarrow \hat{\mathbb{G}}_a$  of  $\mathbb{Z}$ -module schemes over  $\mathbb{Z}_p$ , where the  $\mathbb{Z}$ -module scheme  $E$  is an elliptic curve over  $\mathbb{Z}_p$  and  $J^n E$  is its  $p$ -typical arithmetic jet space defined above, we will look at, for example, characters  $J^n E \rightarrow \hat{\mathbb{G}}_a$  of  $\mathbb{F}_q[t]$ -module schemes over  $\mathbb{F}_q[[t]]$ , where  $E$  is a Drinfeld  $\mathbb{F}_q[t]$ -module over  $\mathbb{F}_q[[t]]$  and  $J^n E$  is its function-field arithmetic jet space—in other words, the Greenberg transform but with “ $t$ -typical” Witt vectors. The most important result in this paper is the construction of a canonical  $F$ -crystal  $\mathbf{H}(E)$  which comes with a Hodge-type filtration and a morphism  $\mathbf{H}(E) \rightarrow \mathbf{H}_{\text{dR}}(E)$  to the usual de Rham cohomology preserving the filtration. As a consequence of the methods that go into the construction of  $\mathbf{H}(E)$ , we also establish a stronger integral version of Buium's result that  $\mathbf{X}_\infty(E)$  itself is freely generated by a single element as an  $R\{\phi^*\}$ -module. Here, we would like to emphasize that all the fundamental principles that go into our approach also work for  $p$ -adic elliptic curves.

Before we describe our main results in detail, we wish to fix a few notations. Let  $\mathbb{F}_q$  be the finite field with  $q$  elements and  $A$  is the coordinate ring of  $X \setminus \{\infty\}$ , where  $X$  is a projective, geometrically connected, smooth curve over  $\mathbb{F}_q$  and  $\infty$  a  $\mathbb{F}_q$ -point on it. Let  $\mathfrak{p}$  be a fixed maximal ideal of  $A$ , and let  $\pi$  be an element of  $\mathfrak{p} \setminus \mathfrak{p}^2$ . Let  $R$  be an  $A$ -algebra which is a complete discrete valuation ring with maximal ideal  $\pi R$  and which has a lift  $\phi: R \rightarrow R$  of the  $q$ -power Frobenius from  $R/\pi R$ , where

$\hat{q} = |A/\mathfrak{p}|$ . Then one can consider the operator on  $R$  given by  $\delta x = \frac{\phi(x) - x^{\hat{q}}}{\pi}$ . It is called the  $\pi$ -derivation associated to  $\phi$ .

Then as in the mixed-characteristic case above, one can define the  $t$ -typical Witt vectors and hence the  $t$ -typical arithmetic jet space functor. For any (formal)  $A$ -module scheme  $E$  over  $R$ , the jet space also  $J^n E$  has a natural (formal)  $A$ -module-scheme structure. However, we would like to remark here that for all  $n \geq 1$ , the  $J^n E$  are not Anderson modules. Then we let  $\mathbf{X}_n(E)$  denote the set of  $A$ -linear differential characters of order  $n$ , that is, the set of homomorphisms  $J^n E \rightarrow \hat{\mathbb{G}}_a$  of (formal)  $A$ -module schemes over  $R$ . Finally, we form their direct limit  $\mathbf{X}_\infty(E)$ , which is naturally an  $R\{\phi^*\}$ -module, as above.

We say  $E$  splits at  $m$  if  $\mathbf{X}_m(E) \neq \{0\}$  but  $\mathbf{X}_i(E) = \{0\}$  for all  $0 \leq i \leq m - 1$ . Then we show that  $m$  satisfies  $1 \leq m \leq r$ , where  $r$  is the rank of  $E$ , and that  $\mathbf{X}_m(E)$  is a free  $R$ -module with a canonical basis element  $\Theta_m \in \mathbf{X}_m(E)$ , depending only on our chosen coordinate on  $E$ . In the case when the rank  $r$  is 2, we have  $m = 2$  unless  $E$  admits a lift of Frobenius compatible with the  $A$ -module structure on  $E$ , in which case  $m = 1$ . Then our first main theorem is a strengthened version of Buium's result in [Bui2].

**Theorem 1.1.** *Let  $E$  be a Drinfeld module that splits at  $m$ . Then the  $R$ -module  $\mathbf{X}_m(E)$  is free of rank 1, and it freely generates  $\mathbf{X}_\infty(E)$  as an  $R\{\phi^*\}$ -module in the sense that the canonical map  $R\{\phi^*\} \otimes_R \mathbf{X}_m(E) \rightarrow \mathbf{X}_\infty(E)$  is an isomorphism.*

Let us now proceed to our second result. Let  $u : J^n E \rightarrow E$  be the usual projection map and put  $N^n = \ker u$ . Since  $u$  is  $A$ -linear,  $N^n$  is a formal  $A$ -module scheme of relative dimension  $n$  over  $\mathrm{Spf} R$ . For each  $n \geq 1$ , we show in proposition 7.2 that there is a lift of Frobenius  $\mathfrak{f} : N^{n+1} \rightarrow N^n$  making the system  $\{N^n\}$  into a prolongation sequence with respect the obvious projection map  $u : N^{n+1} \rightarrow N^n$ . We call  $\mathfrak{f}$  the *lateral Frobenius*. However,  $\mathfrak{f}$  is not compatible with  $i$  and  $\phi : J^{n+1} E \rightarrow J^n E$  in the obvious way, that is, it is not true that  $\phi \circ i = i \circ \mathfrak{f}$  holds. In fact, we can not expect it to be true because that would induce an  $A$ -linear lift of Frobenius on  $E$  which is not the case to start with. Instead we have

$$\phi^2 \circ i = \phi \circ i \circ \mathfrak{f}.$$

In section 9, we construct a canonical  $F$ -crystal attached to  $E$ . The  $F$ -crystal, denoted  $\mathbf{H}(E)$ , is an  $R$ -module which has a semi-linear operator  $\mathfrak{f}^*$  (induced from  $\mathfrak{f}$ ) on it and is of rank  $m$ , which we emphasize can be strictly smaller than  $r$ . The module  $\mathbf{H}(E)$  also has a Hodge-type filtration and canonically maps to the de Rham cohomology of  $E$ , with its Hodge filtration.

**Theorem 1.2.** *There is a canonical map between exact sequences*

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathbf{X}_m(E) & \longrightarrow & \mathbf{H}(E) & \longrightarrow & \mathbf{I}(E) \longrightarrow 0 \\ & & \downarrow \Upsilon & & \downarrow \Phi & & \downarrow \wr \\ 0 & \longrightarrow & \mathrm{Lie}(E)^* & \longrightarrow & \mathbf{H}_{\mathrm{dR}}(E) & \longrightarrow & \mathrm{Ext}(E, \hat{\mathbb{G}}_a) \longrightarrow 0 \end{array}$$

*Moreover, the operator  $\mathfrak{f}^*$  on  $\mathbf{H}(E)$  descends to its image under  $\Phi$ .*

There is a close connection between these two theorems—in fact, our proof of theorem 1.1 goes by way of theorem 1.2.

Finally, we conclude the paper with some explicit computations of the structure constants of the  $F$ -crystal  $\mathbf{H}(E)$ , which are new differential modular forms.

To a Drinfeld module  $E$ , the crystalline theory also attaches an  $F$ -crystal  $\mathbf{H}_{\text{crys}}(E)$ . It appears that our  $\mathbf{H}(E)$  has subtle connections with  $\mathbf{H}_{\text{crys}}(E)$ , but it also appears that any such connection would be indirect. This is because  $\mathbf{H}(E)$ , unlike  $\mathbf{H}_{\text{crys}}(E)$ , has a fundamentally differential-algebraic nature in that it lies not over a point of the moduli space of Drinfeld modules but over a point of the jet space of the moduli space. For instance, the computations in section 10 show the structure constants of  $\mathbf{H}(E)$  do involve the higher  $\pi$ -derivatives of the structure constants of the Drinfeld module. It would be interesting to understand the exact nature of the relationship between  $\mathbf{H}(E)$  and the crystalline cohomology groups, as well as the étale cohomology groups and the other constructions in  $\pi$ -adic Hodge theory. This is all the more true because, as we remarked before, the techniques developed in this paper have analogues for  $p$ -adic elliptic curves, and as a result, we do obtain an analogous construction of the  $F$ -crystal  $\mathbf{H}(E)$  for elliptic curves.

## 2. NOTATION

Let us fix some notation which will hold throughout the paper. Let  $q = p^h$  where  $p$  is a prime and  $h \geq 1$ . Let  $X$  be a projective, geometrically connected, smooth curve over  $\mathbb{F}_q$ . Fix an  $\mathbb{F}_q$ -rational point  $\infty$  on  $X$ . Let  $A$  denote the Dedekind domain  $\mathcal{O}(X \setminus \{\infty\})$ . Let  $\mathfrak{p}$  be a maximal ideal of  $A$ , and let  $\hat{A}$  denote the  $\mathfrak{p}$ -adic completion of  $A$ . Let  $t$  be an element of  $\mathfrak{p} \setminus \mathfrak{p}^2$ , and let  $\pi$  denote its image in  $\hat{A}$ . Then  $\pi$  generates the maximal ideal  $\hat{\mathfrak{p}}$  of  $\hat{A}$ . Let  $k$  denote the residue field  $A/\mathfrak{p}$ , and let  $\hat{q}$  denote its cardinality. Note that the quotient map  $\hat{A} \rightarrow k$  has a unique section. Thus  $\hat{A}$  is not just an  $\mathbb{F}_q$ -algebra but also canonically a  $k$ -algebra.

Now let  $R$  be an  $\hat{A}$ -algebra which is  $\mathfrak{p}$ -adically complete and flat, or equivalently  $\pi$ -torsion free. Thus the composition  $\theta : A \rightarrow \hat{A} \rightarrow R$  is injective (assuming  $R \neq \{0\}$ ) and hence one says that  $\theta$  is of *generic characteristic*. Let us also fix an  $\hat{A}$ -algebra endomorphism  $\phi : R \rightarrow R$  which lifts the  $\hat{q}$ -power Frobenius modulo  $\mathfrak{p}R$ :

$$\phi(x) \equiv x^{\hat{q}} \pmod{\mathfrak{p}R}.$$

Do note that the identity map on  $\hat{A}$  does indeed lift the  $\hat{q}$ -power Frobenius on  $\hat{A}/\hat{\mathfrak{p}}$ .

Also note that not all rings  $R$  admit such a Frobenius lift; so the existence of  $\phi$  does place a restriction on  $R$ . For our main results,  $R$  will in the end be a discrete valuation ring, most importantly the completion of the maximal unramified extension of  $\hat{A}$ . So the reader may assume this from the start. But some form of our results should hold in general, and with essentially the same proofs. This is of some interest, for instance when  $R$  is the coordinate ring of the moduli space of Drinfeld modules of a given rank. With an eye to the future, we have not assumed that  $R$  is a discrete valuation ring where it is easily avoided, in sections 3–7.

Let  $K$  denote  $R[1/\pi]$ , and for any  $R$ -module  $M$  write  $M_K = K \otimes_R M$ . Finally, let  $S$  denote  $\text{Spf } R$ .

## 3. FUNCTION-FIELD WITT VECTORS

Witt vectors over Dedekind domains with finite residue fields were introduced in [Bo1]. We will only work over  $\hat{A}$ , which is the ring of integers of a local field of characteristic  $p$ , and here they were introduced earlier in [D76]. The basic results can be developed exactly as in any of the usual developments of the  $p$ -typical Witt vectors. The only difference is that in all formulas any  $p$  in a coefficient is replaced with a  $\pi$  and any  $p$  in an exponent is replaced with a  $\hat{q}$ .

**3.1. Frobenius lifts and  $\pi$ -derivations.** Let  $B$  be an  $R$ -algebra, and let  $C$  be a  $B$ -algebra with structure map  $u : B \rightarrow C$ . In this paper, a ring homomorphism  $\psi : B \rightarrow C$  will be called a *lift of Frobenius* (relative to  $u$ ) if it satisfies the following:

- (1) The reduction mod  $\pi$  of  $\psi$  is the  $\hat{q}$ -power Frobenius relative to  $u$ , that is,  $\psi(x) \equiv u(x)^{\hat{q}} \pmod{\pi C}$ .
- (2) The restriction of  $\psi$  to  $R$  coincides with the fixed  $\phi$  on  $R$ , that is, the following diagram commutes

$$\begin{array}{ccc} B & \xrightarrow{\psi} & C \\ \uparrow & & \uparrow \\ R & \xrightarrow{\phi} & R \end{array}$$

A  $\pi$ -*derivation*  $\delta$  from  $B$  to  $C$  means a set-theoretic map  $\delta : B \rightarrow C$  satisfying the following for all  $x, y \in B$

$$\begin{aligned} \delta(x + y) &= \delta(x) + \delta(y) \\ \delta(xy) &= u(x)^{\hat{q}}\delta(y) + \delta(x)u(y)^{\hat{q}} + \pi\delta(x)\delta(y) \end{aligned}$$

such that for all  $r \in R$ , we have

$$\delta(r) = \frac{\phi(r) - r^{\hat{q}}}{\pi}.$$

When  $C = B$  and  $u$  is the identity map, we will call this simply a  $\pi$ -derivation on  $B$ .

It follows that the map  $\phi : B \rightarrow C$  defined as

$$\phi(x) := u(x)^{\hat{q}} + \pi\delta(x)$$

is a lift of Frobenius in the sense above. On the other hand, for any flat  $R$ -algebra  $B$  with a lift of Frobenius  $\phi$ , one can define the  $\pi$ -derivation  $\delta(x) = \frac{\phi(x) - x^{\hat{q}}}{\pi}$  for all  $x \in B$ .

Note that this definition depends on the choice of uniformizer  $\pi$ , but in a transparent way: if  $\pi'$  is another uniformizer, then  $\delta(x)\pi/\pi'$  is a  $\pi'$ -derivation. This correspondence induces a bijection between  $\pi$ -derivations  $B \rightarrow C$  and  $\pi'$ -derivations  $B \rightarrow C$ .

**3.2. Witt vectors.** We will present three different points of view on function-field Witt vectors, all parallel to the mixed characteristic case. But there is perhaps one unfamiliar element below, which is that we will work relative to our general base  $R$ , and it already has a lift of Frobenius. The consequence is that we need to pay attention to certain twists of the scalars by Frobenius, which are invisible over

the absolute base  $R = \hat{A}$ . However this unfamiliar element has nothing to do with the difference between mixed and equal characteristic and only with the difference between the relative and the absolute setting.

Let  $B$  be an  $R$ -algebra with structure map  $u : R \rightarrow B$ .

(1) The ring  $W(B)$  of  $\pi$ -typical Witt vectors can be defined as the unique (up to unique isomorphism)  $R$ -algebra  $W(B)$  with a  $\pi$ -derivation  $\delta$  on  $W(B)$  and an  $R$ -algebra homomorphism  $W(B) \rightarrow B$  such that, given any  $R$ -algebra  $C$  with a  $\pi$ -derivation  $\delta$  on it and an  $R$ -algebra map  $f : C \rightarrow B$ , there exists a unique  $R$ -algebra homomorphism  $g : C \rightarrow W(B)$  such that the diagram

$$\begin{array}{ccc} W(B) & & \\ \downarrow & \swarrow g & \\ B & \xleftarrow{f} & C \end{array}$$

commutes and  $g \circ \delta = \delta \circ g$ . Thus  $W$  is the right adjoint of the forgetful functor from  $R$ -algebras with  $\pi$ -derivation to  $R$ -algebras. For details, see section 1 of [Bo1]. This approach follows that of [Jo] to the usual  $p$ -typical Witt vectors.

(2) If we restrict to flat  $R$ -algebras  $B$ , then we can ignore the concept of  $\pi$ -derivation and define  $W(B)$  simply by expressing the universal property above in terms of Frobenius lifts, as follows. Given a flat  $R$ -algebra  $B$ , the ring  $W(B)$  is the unique (up to unique isomorphism) flat  $R$ -algebra  $W(B)$  with a lift of Frobenius (in the sense above)  $F : W(B) \rightarrow W(B)$  and an  $R$ -algebra homomorphism  $W(B) \rightarrow B$  such that for any flat  $R$ -algebra  $C$  with a lift of Frobenius  $\phi$  on it and an  $R$ -algebra map  $f : C \rightarrow B$ , there exists a unique  $R$ -algebra homomorphism  $g : C \rightarrow W(B)$  such that the diagram

$$\begin{array}{ccc} W(B) & & \\ \downarrow & \swarrow g & \\ B & \xleftarrow{f} & C \end{array}$$

commutes and  $g \circ \phi = F \circ g$ .

(3) Finally, one can also define Witt vectors in terms of the Witt polynomials. For each  $n \geq 0$  let us define  $B^{\phi^n}$  to be the  $R$ -algebra with structure map  $R \xrightarrow{\phi^n} R \xrightarrow{u} B$  and define the *ghost rings* to be the product  $R$ -algebras  $\Pi_{\phi}^n B = B \times B^{\phi} \times \cdots \times B^{\phi^n}$  and  $\Pi_{\phi}^{\infty} B = B \times B^{\phi} \times \cdots$ . Then for all  $n \geq 1$  there exists a *restriction*, or *truncation*, map  $T_w : \Pi_{\phi}^n B \rightarrow \Pi_{\phi}^{n-1} B$  given by  $T_w(w_0, \dots, w_n) = (w_0, \dots, w_{n-1})$ . We also have the left shift *Frobenius* operators  $F_w : \Pi_{\phi}^n B \rightarrow \Pi_{\phi}^{n-1} B$  given by  $F_w(w_0, \dots, w_n) = (w_1, \dots, w_n)$ . Note that  $T_w$  is an  $R$ -algebra morphism, but  $F_w$  lies over the Frobenius endomorphism  $\phi$  of  $R$ .

Now as sets define

$$(3.1) \quad W_n(B) = B^{n+1},$$

and define the set map  $w : W_n(B) \rightarrow \Pi_{\phi}^n B$  by  $w(x_0, \dots, x_n) = (w_0, \dots, w_n)$  where

$$(3.2) \quad w_i = x_0^{\hat{q}^i} + \pi x_1^{\hat{q}^{i-1}} + \cdots + \pi^i x_i$$

are the *Witt polynomials*. The map  $w$  is known as the *ghost* map. (Do note that under the traditional indexing our  $W_n$  would be denoted  $W_{n+1}$ .) We can

then define the ring  $W_n(B)$ , the ring of truncated  $\pi$ -typical Witt vectors, by the following theorem as in the  $p$ -typical case [H05]:

**Theorem 3.1.** *For each  $n \geq 0$ , there exists a unique functorial  $R$ -algebra structure on  $W_n(B)$  such that  $w$  becomes a natural transformation of functors of  $R$ -algebras.*

Note that, unlike with the usual Witt vectors in mixed characteristic, addition for function-field Witt vectors is performed componentwise. This is because the Witt polynomials (3.2) are additive. This might appear to defeat the whole point of Witt vectors and arithmetic jet spaces. But this is not so. The reason is that while the additive structure is the componentwise one, the  $A$ -module structure is not. So the difference is only that, unlike in mixed characteristic where  $A = \mathbb{Z}$ , a group structure is weaker than  $A$ -module structure. In fact, because the Witt polynomials are  $k$ -linear, the  $k$ -vector space structure on  $W_n(B)$  is the componentwise one. This is just like with the  $p$ -typical Witt vectors, where multiplication by roots of  $x^p - x$  can be performed componentwise.

**3.3. Operations on Witt vectors.** Now we recall some important operators on the Witt vectors. There are the *restriction*, or *truncation*, maps  $T : W_n(B) \rightarrow W_{n-1}(B)$  given by  $T(x_0, \dots, x_n) = (x_0, \dots, x_{n-1})$ . Note that  $W(B) = \lim_{\leftarrow} W_n(B)$ . There is also the *Frobenius* ring homomorphism  $F : W_n(B) \rightarrow W_{n-1}(B)$ , which can be described in terms of the ghost map. It is the unique map which is functorial in  $B$  and makes the following diagram commutative

$$(3.3) \quad \begin{array}{ccc} W_n(B) & \xrightarrow{w} & \Pi_{\phi}^n B \\ F \downarrow & & \downarrow F_w \\ W_{n-1}(B) & \xrightarrow{w} & \Pi_{\phi}^{n-1} B^n \end{array}$$

As with the ghost components,  $T$  is an  $R$ -algebra map but  $F$  lies over the Frobenius endomorphism  $\phi$  of  $R$ .

Next we have the *Verschiebung*  $V : W_{n-1}(B) \rightarrow W_n(B)$  given by

$$V(x_0, \dots, x_{n-1}) = (0, x_0, \dots, x_{n-1}).$$

Let  $V_w : \Pi_{\phi}^{n-1} B \rightarrow \Pi_{\phi}^n B$  be the additive map given by

$$V_w(w_0, \dots, w_{n-1}) = (0, \pi w_0, \dots, \pi w_{n-1}).$$

Then the Verschiebung  $V$  makes the following diagram commute:

$$(3.4) \quad \begin{array}{ccc} W_{n-1}(B) & \xrightarrow{w} & \Pi_{\phi}^{n-1} B \\ V \downarrow & & \downarrow V_w \\ W_n(B) & \xrightarrow{w} & \Pi_{\phi}^n B \end{array}$$

For all  $n \geq 0$  the Frobenius and the Verschiebung satisfy the identity

$$(3.5) \quad FV(x) = \pi x.$$

The Verschiebung is not a ring homomorphism, but it is  $k$ -linear.

Finally, we have the multiplicative Teichmüller map  $[ ] : B \rightarrow W_n(B)$  given by  $x \mapsto [x] = (x, 0, 0, \dots)$ . Here in the function-field setting,  $[ ]$  is additive and even a homomorphism of  $k$ -algebras.

**3.4. Computing the universal map to Witt vectors.** Given an  $R$ -algebra  $C$  with a  $\pi$ -derivation  $\delta$  and an  $R$ -algebra map  $f : C \rightarrow B$ , we will now describe the universal lift  $g : C \rightarrow W(B)$ . The explicit description of  $g$  leads us to proposition 3.2 which is used in section 10 in computations for Drinfeld modules of rank 2. The reader may skip this subsection without breaking continuity till then.

It is enough to work in the case where both  $B$  and  $C$  are flat over  $R$ . Then the ghost map  $w : W(B) \rightarrow \Pi_\phi^\infty B$  is injective. Consider the map  $[\phi] : C \rightarrow \Pi_\phi^\infty C$  given by  $x \mapsto (x, \phi(x), \phi^2(x), \dots)$ . Then we have the following commutative diagram:

$$\begin{array}{ccccc}
 & & & & C \\
 & & & \nearrow g & \downarrow [\phi] \\
 & & & f \circ [\phi] & \\
 W(B) & \xrightarrow{w} & \Pi_\phi^\infty B & \xleftarrow{f} & \Pi_\phi^\infty C \\
 \downarrow F & & \downarrow F_w & & \downarrow F_w \\
 W(B) & \xrightarrow{w} & \Pi_\phi^\infty B & \xleftarrow{f} & \Pi_\phi^\infty C
 \end{array}$$

Thus the map  $f \circ [\phi] : C \rightarrow \Pi_\phi^\infty B$  factors through  $W(B)$  as our universal map  $g : C \rightarrow W(B)$ .

Let us now give an inductive description of the map  $g$ . Write

$$g(x) = (x_0, x_1, \dots) \in W(B).$$

Then from the above diagram  $w \circ g = f \circ [\phi]$ . Therefore the vector  $(x_0, x_1, \dots)$  is the unique solution to the system of equations

$$(3.6) \quad x_0^{\hat{q}^n} + \pi x_1^{\hat{q}^{n-1}} + \dots + \pi^n x_n = f(\phi^n(x)),$$

for  $n \geq 0$ . For example, we have  $x_0 = f(x)$  and  $x_1 = f(\delta(x))$ .

Now consider the case where  $B$  itself has a  $\pi$ -derivation,  $C = B$ , and  $f = \mathbb{1}$ . For any  $x \in B$ , let us write  $x^{(n)} := \delta^n(x)$ , or simply  $x' = \delta(x)$ ,  $x'' = \delta^2(x)$  and so on.

**Proposition 3.2.** *We have  $x_0 = x$ ,  $x_1 = x'$  and  $x_2 = x'' + \pi^{\hat{q}-2}(x')^{\hat{q}}$ .*

*Proof.* As stated above, equalities  $x_0 = x$  and  $x_1 = x'$  follow immediately from (3.6). For  $n = 2$ , we have

$$\begin{aligned}
 x_0^{\hat{q}^2} + \pi x_1^{\hat{q}} + \pi^2 x_2 &= \phi^2(x) \\
 &= \phi(x^{\hat{q}} + \pi x') \\
 &= \phi(x)^{\hat{q}} + \pi \phi(x') \\
 &= x^{\hat{q}^2} + \pi^{\hat{q}}(x')^{\hat{q}} + \pi((x')^{\hat{q}} + \pi x'')
 \end{aligned}$$

And therefore we have  $x_2 = x'' + \pi^{\hat{q}-2}(x')^{\hat{q}}$ .  $\square$

#### 4. $A$ -MODULE SCHEMES, JET SPACES AND PRELIMINERIES

An  $A$ -module scheme over  $S = \mathrm{Spf} R$  is by definition a pair  $(E, \varphi_E)$ , where  $E$  is a commutative group object in the category of  $S$ -schemes and  $\varphi_E : A \rightarrow \mathrm{End}(E/S)$  is a ring map. (Here and below, by a scheme over the formal scheme  $S$ , we mean a formal scheme formed from a compatible family of schemes over the schemes  $\mathrm{Spec} R/\mathfrak{p}^n R$ .) Then the tangent space  $T_0 E$  at the identity has two  $A$ -modules



structures: one coming by restriction of the usual  $R$ -module structure to  $A$ , and the other coming from differentiating  $\varphi_E$ . We will say that  $(E, \varphi_E)$  is *strict* if these two  $A$ -module structures coincide, and *admissible* if it is both strict and isomorphic to the additive group  $\hat{\mathbb{G}}_a = \hat{\mathbb{G}}_{a/S}$  as a group scheme. We will denote this induced map to tangent space as  $\theta : A \rightarrow R$ . (Note that it is best practice to require only the isomorphism with  $\hat{\mathbb{G}}_a$  to exist locally on  $S$ . So below, our Drinfeld modules would more properly be called *coordinatized* Drinfeld modules.)

A *Drinfeld module*  $(E, \varphi_E)$  of rank  $r$  is an admissible  $A$ -module scheme over  $S$  such that for each non-zero  $a \in A$ , the group scheme  $\ker(\varphi_E(a))$  is finite of degree  $|a|^r = q^{-r \text{ord}_\infty(a)}$  over  $S$ .

**Proposition 4.1.** *If  $f$  is an endomorphism of the  $\mathbb{F}_q$ -module scheme  $\hat{\mathbb{G}}_{a/S}$  over  $S$ , then it is of the form*

$$f(x) = \sum_{i=0}^{\infty} a_i x^{q^i},$$

where  $f$  is a restricted power series, meaning  $a_i \rightarrow 0$   $\pi$ -adically as  $i \rightarrow \infty$ .

*Proof.* Let  $f \in \text{Hom}(\hat{\mathbb{G}}_a, \hat{\mathbb{G}}_a)$  be an additive endomorphism of  $\hat{\mathbb{G}}_a$ . Then  $f$  is given a restricted power series  $\sum_i b_i x^i$  such that  $b_i \rightarrow 0$  as  $i \rightarrow \infty$ . Since  $f$  is additive, we have  $b_i = 0$  unless  $i$  is a power of  $p$ . Second, because  $f$  is  $\mathbb{F}_q$ -linear, we have  $\sum_i b_{p^i} (cx)^{p^i} = c \sum_i b_{p^i} x^{p^i}$  for all  $c \in \mathbb{F}_q$ . Considering the case where  $c$  is a generator of  $\mathbb{F}_q^*$ , we see this implies  $b_{p^i} = 0$  unless  $p^i$  is a power of  $q$ .  $\square$

Let  $R\{\tau\}^\wedge$  be the subring of  $R\{\{\tau\}\}$  consisting of (twisted) restricted power series. Then by proposition 4.1, the  $\mathbb{F}_q$ -linear morphisms between two admissible  $A$ -module schemes  $E_1$  and  $E_2$  over  $\text{Spf } R$  are given in coordinates by elements in  $R\{\tau\}^\wedge$  where  $\tau$  acts as  $\tau(x) = x^q$ :

$$(4.1) \quad \text{Hom}_{\mathbb{F}_q}(E_1, E_2) = R\{\tau\}^\wedge.$$

**4.1. Prolongation sequences and jet spaces.** Let  $X$  and  $Y$  be schemes over  $S = \text{Spf } R$ . We say a pair  $(u, \delta)$  is a *prolongation*, and write  $Y \xrightarrow{(u, \delta)} X$ , if  $u : Y \rightarrow X$  is a map of schemes over  $S$  and  $\delta : \mathcal{O}_X \rightarrow u_* \mathcal{O}_Y$  is a  $\pi$ -derivation making the following diagram commute:

$$\begin{array}{ccc} R & \longrightarrow & u_* \mathcal{O}_Y \\ \delta \uparrow & & \uparrow \delta \\ R & \longrightarrow & \mathcal{O}_X \end{array}$$

Following [Bui3], a *prolongation sequence* is a sequence of prolongations

$$\text{Spf } R \xleftarrow{(u, \delta)} T^0 \xleftarrow{(u, \delta)} T^1 \xleftarrow{(u, \delta)} \dots,$$

where each  $T^n$  is a scheme over  $S$ . We will often use the notation  $T^*$  or  $\{T_n\}_{n \geq 0}$ . Note that if the  $T^n$  are flat over  $\text{Spf } R$  then having a  $\pi$ -derivation  $\delta$  is equivalent to having lifts of Frobenius  $\phi : T^{n+1} \rightarrow T^n$ .

Prolongation sequences form a category  $\mathcal{C}_{S^*}$ , where a morphism  $f : T^* \rightarrow U^*$  is a family of morphisms  $f^n : T^n \rightarrow U^n$  commuting with both the  $u$  and  $\delta$ , in the evident sense. This category has a final object  $S^*$  given by  $S^n = \text{Spf } R$  for all  $n$ , where each  $u$  is the identity and each  $\delta$  is the given  $\pi$ -derivation on  $R$ .

For any scheme  $Y$  over  $S$ , for all  $n \geq 0$  we define the  $n$ -th jet space  $J^n X$  (relative to  $S$ ) as

$$J^n X(Y) := \mathrm{Hom}_S(W_n^*(Y), X)$$

where  $W_n^*(Y)$  is defined as in [Bo2]. We will not define  $W_n^*(Y)$  in full generality here. Instead, we will define  $\mathrm{Hom}_S(W_n^*(Y), X)$  in the affine case, and that will be sufficient for the purposes of this paper. Write  $X = \mathrm{Spf} A$  and  $Y = \mathrm{Spf} B$ . Then  $W_n^*(Y) = \mathrm{Spf} W_n(B)$  and  $\mathrm{Hom}_S(W_n^*Y, X)$  is  $\mathrm{Hom}_R(A, W_n(B))$ , the set of  $R$ -algebra homomorphisms  $A \rightarrow W_n(B)$ .

Then  $J^*X := \{J^n X\}_{n \geq 0}$  forms a prolongation sequence and is called the *canonical prolongation sequence* [Bui3]. By [Bui3], [Bo2],  $J^*X$  satisfies the following universal property—for any  $T^* \in \mathcal{C}_{S^*}$  and  $X$  a scheme over  $S^0$ , we have

$$\mathrm{Hom}(T^0, X) = \mathrm{Hom}_{\mathcal{C}_{S^*}}(T^*, J^*X)$$

Let  $X$  be a scheme over  $S = \mathrm{Spf} R$ . Define  $X^{\phi^n}$  by  $X^{\phi^n}(B) := X(B^{\phi^n})$  for any  $R$ -algebra  $B$ . In other words,  $X^{\phi^n}$  is  $X \times_{S, \phi^n} S$ , the pull-back of  $X$  under the map  $\phi^n : S \rightarrow S$ . Next define

$$\Pi_\phi^n X = X \times_S X^\phi \times_S \cdots \times_S X^{\phi^n}.$$

Then for any  $R$ -algebra  $B$  we have  $X(\Pi_\phi^n B) = X(B) \times_S \cdots \times_S X^{\phi^n}(B)$ . Thus the ghost map  $w$  in theorem 3.1 defines a map of  $S$ -schemes

$$w : J^n X \rightarrow \Pi_\phi^n X.$$

Note that  $w$  is injective when evaluated on points with coordinates in any flat  $R$ -algebra.

The operators  $F$  and  $F_w$  in (3.3) induce maps  $\phi$  and  $\phi_w$  as follows

$$(4.2) \quad \begin{array}{ccc} J^n X & \xrightarrow{w} & \Pi_\phi^n X \\ \phi \downarrow & & \downarrow \phi_w \\ J^{n-1} X & \xrightarrow{w} & \Pi_\phi^{n-1} X \end{array}$$

where  $\phi_w$  is the left-shift operator given by

$$\phi_w(w_0, \dots, w_n) = (\phi_S(w_1), \dots, \phi_S(w_n)),$$

and where  $\phi_S : X^{\phi^i} \rightarrow X^{\phi^{i-1}}$  is the composition given in the following diagram:

$$(4.3) \quad \begin{array}{ccccc} X^{\phi^i} & \xrightarrow{\sim} & X^{\phi^{i-1}} \times_{S, \phi} S & \longrightarrow & X^{\phi^{i-1}} \\ & & \downarrow & & \downarrow \\ & & S & \xrightarrow{\phi} & S. \end{array}$$

Now let  $E$  be an  $A$ -module scheme over  $S$  with action map  $A \xrightarrow{\varphi_E} \mathrm{End}_S(E)$ . Then the functor it represents takes values in  $A$ -modules, and hence so does the functor  $B \mapsto E(W_n(B))$ . In this way, for each  $n \geq 0$ , the  $S$ -scheme  $J^n E$  comes with an  $A$ -module structure. We denote it by  $\varphi_{J^n E} : A \rightarrow \mathrm{End}_S(J^n E)$ . Similarly,  $\varphi_E$  induces an  $A$ -linear structure  $\varphi_{E^{\phi^n}}$  on each  $E^{\phi^n}$ . In this case, it is easy to describe explicitly. It is the componentwise one:

$$\varphi_{\Pi_\phi^n E}(w_0, \dots, w_n) = (\varphi_E(w_0), \dots, \varphi_{E^{\phi^n}}(w_n)).$$

The ghost map  $w : J^n E \rightarrow \Pi_\phi^n E$  and the truncation map  $u : J^n E \rightarrow J^{n-1} E$  homomorphisms of  $A$ -module schemes over  $S$ . This is because they are given by applying the  $A$ -module scheme  $E$  to the  $R$ -algebra maps  $w : W_n(B) \rightarrow \Pi_\phi^n B$  and  $T : W_n(B) \rightarrow W_{n-1}(B)$ . On the other hand, the Frobenius map  $\phi : J^n E \rightarrow J^{n-1} E$  is a homomorphisms of  $A$ -module schemes lying over the Frobenius endomorphism  $\phi$  of  $S$ . In other words, the induced map  $J^n E \rightarrow (J^{n-1} E)^\phi$  is a homomorphism of  $A$ -module schemes over  $S$ .

**4.2. Coordinates on jet spaces.** Given an isomorphism of  $S$ -schemes  $E \rightarrow \hat{\mathbb{G}}_a$ , we can identify  $(J^n E)(B)$  with  $W_n(B)$  and hence, using (3.1), with  $B^{n+1}$ . In particular, given a coordinate  $x$  on an admissible  $A$ -module scheme  $E$ , this identification provides a canonical system of coordinates  $(x_0, \dots, x_n)$  on  $J^n E$ . We will use these *Witt coordinates* without further comment. We emphasize once again that there are other canonical systems of coordinates on  $J^n E$ , for instance the *Buium–Joyal* coordinates denoted  $x, x', x'', \dots$ . They are related by the formulas of proposition 3.2. Each has their own advantages.

**4.3. Character groups.** Given a prolongation sequence  $T^*$  we can define its shift  $T^{*+n}$  by  $(T^{*+n})^j := T^{n+j}$  for all  $j$  [Bui3].

$$\mathrm{Spf} R \xleftarrow{(u,\delta)} T^n \xleftarrow{(u,\delta)} T^{n+1} \dots$$

We define a  $\delta$ -morphism of order  $n$  from  $X$  to  $Y$  to be a morphism  $J^{*+n} X \rightarrow J^* Y$  of prolongation sequences. We define a *character of order  $n$* ,  $\Theta : (E, \varphi_E) \rightarrow (\hat{\mathbb{G}}_a, \varphi_{\hat{\mathbb{G}}_a})$  to be a  $\delta$ -morphism of order  $n$  from  $E$  to  $\hat{\mathbb{G}}_a$  which is also a homomorphism of  $A$ -module objects. By the universal property of jet schemes [Bui3], an order  $n$  character is equivalent to a homomorphism  $\Theta : J^n E \rightarrow \hat{\mathbb{G}}_a$  of  $A$ -module schemes over  $S$ . We denote the group of characters of order  $n$  by  $\mathbf{X}_n(E)$ . So we have

$$\mathbf{X}_n(E) = \mathrm{Hom}_A(J^n E, \hat{\mathbb{G}}_a),$$

which one could take as an alternative definition. Note that  $\mathbf{X}_n(E)$  comes with an  $R$ -module structure since  $\hat{\mathbb{G}}_a$  is an  $R$ -module scheme over  $S$ . Also the inverse system  $J^{n+1} E \xrightarrow{u} J^n E$  defines a directed system

$$\mathbf{X}_n(E) \xrightarrow{u^*} \mathbf{X}_{n+1}(E) \xrightarrow{u^*} \dots$$

via pull back. Each morphism  $u^*$  is injective because each  $u$  has a section (typically not  $A$ -linear). We then define  $\mathbf{X}_\infty(E)$  to be the direct limit  $\lim_n \mathbf{X}_n(E)$ .

Similarly, pre-composing with the Frobenius map  $\phi : J^{n+1} E \rightarrow J^n E$  induces a Frobenius operator  $\phi : \mathbf{X}^n(E) \rightarrow \mathbf{X}^{n+1}(E)$ . However since  $\phi : J^{n+1} E \rightarrow J^n E$  is not a morphism over  $\mathrm{Spf} R$  but instead lies over the Frobenius endomorphism  $\phi$  of  $S$ , some care is required. Consider the relative Frobenius morphism  $\phi_R$ , defined

to be the unique morphism making the following diagram commute:

$$\begin{array}{ccc}
J^{n+1}E & \xrightarrow{\phi} & J^n E \\
\downarrow \phi_R & \searrow \phi & \downarrow \\
J^n E \times_{(\mathrm{Spf} R), \phi} \mathrm{Spf} R & \longrightarrow & J^n E \\
\downarrow & & \downarrow \\
\mathrm{Spf} R & \xrightarrow{\phi} & \mathrm{Spf} R
\end{array}$$

Then  $\phi_R$  is a morphism of  $A$ -module formal schemes over  $\mathrm{Spf} R$ . Now given a  $\delta$ -character  $\Theta : J^n E \rightarrow \hat{\mathbb{G}}_a$ , define  $\phi^* \Theta$  to be the composition

$$(4.4) \quad J^{n+1}E \xrightarrow{\phi_R} J^n E \times_{(\mathrm{Spf} R), \phi} \mathrm{Spf} R \xrightarrow{\Theta \times \mathbb{1}} \hat{\mathbb{G}}_a \times_{(\mathrm{Spf} R), \phi} \mathrm{Spf} R \xrightarrow{\iota} \hat{\mathbb{G}}_a$$

where  $\iota$  is the isomorphism of  $A$ -module schemes over  $S$  coming from the fact that  $\hat{\mathbb{G}}_a$  descends to  $\hat{A}$  as an  $A$ -module scheme. For any  $R$ -algebra  $B$ , the induced morphism on  $B$ -points is

$$E(W_{n+1}(B)) \xrightarrow{E(F)} E(W_n(B)^\phi) \xrightarrow{\Theta_B^\phi} B^\phi \xrightarrow{b \mapsto b} B.$$

Note that this composition  $E(W_{n+1}(B)) \rightarrow B$  is indeed a morphism of  $A$ -modules because identity map  $B^\phi \rightarrow B$  is  $A$ -linear, which is true because  $\phi$  restricted to  $\hat{A}$  is the identity.

Thus we have an additive map  $\mathbf{X}_n(E) \rightarrow \mathbf{X}_{n+1}(E)$  given by  $\Theta \mapsto \phi^* \Theta$ . Note that this map is not  $R$ -linear. However, the map

$$\phi^* : \mathbf{X}_n(E) \longrightarrow \mathbf{X}_{n+1}(E)^\phi, \quad \Theta \mapsto \phi^* \Theta$$

is  $R$ -linear, where  $\mathbf{X}_{n+1}(E)^\phi$  denotes the abelian group  $\mathbf{X}_{n+1}(E)$  with  $R$ -module structure defined by the law  $r \cdot \Theta := \phi(r)\Theta$ . Taking direct limits in  $n$ , we obtain an  $R$ -linear map

$$\mathbf{X}_\infty(E) \longrightarrow \mathbf{X}_\infty(E)^\phi, \quad \Theta \mapsto \phi^* \Theta.$$

In this way,  $\mathbf{X}_\infty(E)$  is a left module over the twisted polynomial ring  $R\{\phi^*\}$  with commutation law  $\phi^* r = \phi(r)\phi^*$ .

## 5. KERNEL OF $u : J^1 E \rightarrow E$

Let  $(E, \varphi_E)$  be an admissible  $A$ -module scheme over  $S = \mathrm{Spf} R$ . By equation (4.1), we can write

$$(5.1) \quad \varphi_E(t) = \sum a_i \tau^i$$

with  $a_i \in R$ ,  $a_i \rightarrow 0$ , and  $a_0 = \pi$ . Let  $N^n$  denote the kernel of the projection  $u : J^n E \rightarrow E$ . Thus we have a short exact sequence of  $A$ -module schemes over  $S$ :

$$0 \rightarrow N^n \rightarrow J^n E \xrightarrow{u} E \rightarrow 0$$

We will show in this section that, when  $q \geq 3$ , there is an isomorphism  $(N^1, \varphi_{N^1}) \rightarrow (\hat{\mathbb{G}}_a, \varphi_{\hat{\mathbb{G}}_a})$  of  $A$ -module schemes, where  $\hat{\mathbb{G}}_a$  denotes the tautological  $A$ -module with the  $A$ -action is given by the usual multiplication of scalars:  $\varphi_{\hat{\mathbb{G}}_a}(a) = a\tau^0$ .

**Proposition 5.1.** *Any  $A$ -linear morphism  $f : E \rightarrow G$  between admissible  $A$ -modules is determined by the induced morphism on tangent spaces. More precisely, if we write  $\varphi_E(t) = \pi\tau^0 + \sum_{j \geq 1} a_j \tau^j$ ,  $\varphi_G(t) = \pi\tau^0 + \sum_{j \geq 1} c_j \tau^j$ , and  $f = \sum_i b_i \tau^i$ , then we have*

$$b_r = (\pi - \pi^{q^r})^{-1} \sum_{i=0}^{r-1} (b_i a_{r-i}^{q^i} - c_{r-i} b_i^{q^{r-i}}).$$

*Proof.* Because  $f$  is  $B$ -linear, we have

$$\left( \sum_{i \geq 0} b_i \tau^i \right) \left( \pi\tau^0 + \sum_{j \geq 1} a_j \tau^j \right) = \left( \pi\tau^0 + \sum_{j \geq 1} c_j \tau^j \right) \left( \sum_{i \geq 0} b_i \tau^i \right).$$

Comparing the coefficients of  $\tau^r$ , we have

$$b_0 a_r^{q^0} + \cdots + b_{r-1} a_1^{q^{r-1}} + b_r \pi^{q^r} = \pi b_r + c_1 b_{r-1}^{q^1} + \cdots + c_r b_0^{q^r}.$$

Therefore we have

$$b_r (\pi - \pi^{q^r}) = \sum_{i=0}^{r-1} (b_i a_{r-i}^{q^i} - c_{r-i} b_i^{q^{r-i}}).$$

Since  $R$  is  $\pi$ -torsion free and  $1 - \pi^{q^r-1}$  is invertible for  $r \geq 1$ , this determines each  $b_r$  uniquely in terms of  $b_0, \dots, b_{r-1}$ . Therefore  $b_0$  determines each  $b_r$ .  $\square$

**Corollary 5.2.** *The  $R$ -module map  $R \rightarrow \text{Hom}_A(\hat{\mathbb{G}}_a, \hat{\mathbb{G}}_a)$  defined by  $b \mapsto b\tau^0$  is an isomorphism.*

Now consider the subset  $S^\dagger \subset R\{\tau\}^\wedge$  defined by

$$(5.2) \quad S^\dagger := \left\{ \sum_{i \geq 0} b_i \tau^i \in R\{\tau\}^\wedge \mid v(b_i) \geq i, \text{ for all } i \text{ and } b_0 \in R^* \right\}.$$

Here, and below, we write  $v(b) \geq i$  to mean simply  $b \in \mathfrak{p}^i R$ .

**Proposition 5.3.**  *$S^\dagger$  is a group under composition.*

*Proof.* The fact that  $S^\dagger$  is a submonoid of  $R\{\tau\}^\wedge$  under composition follows immediately from the law  $b\tau^i \circ c\tau^j = bc^{q^i} \tau^{i+j}$  and linearity. Indeed if  $v(b) \geq i$  and  $v(c) \geq j$ , then  $v(bc^{q^i}) \geq i + j$ .

Now let us show that any element  $f = \sum b_i \tau^i \in S^\dagger$  has an inverse under composition. Let  $g = \sum_{n=0}^{\infty} c_n \tau^n$ , where  $c_0 = b_0^{-1}$  and we define inductively  $c_n = -b_0^{-q^n} (c_0 b_n + c_1 b_{n-1}^{q^1} + \cdots + c_{n-1} b_1^{q^{n-1}})$ . Then it is easy to check that  $g \circ f = \mathbb{1}$ . Take  $n \geq 1$  and assume  $v(c_i) \geq i$  for all  $i = 0, \dots, n-1$ . Then it is enough to show  $v(c_n) \geq n$ . We have  $v(c_n) \geq \min\{v(c_i b_{n-i}^{q^i}) \mid i = 0, \dots, n-1\}$ . Now

$$\begin{aligned} v(c_i b_{n-i}^{q^i}) &= v(c_i) + q^i v(b_{n-i}) \\ &= i + q^i (n - i) \\ &\geq i + (n - i) = n. \end{aligned}$$

Therefore the left inverse  $g$  of  $f$  lies in  $S^\dagger$ .

Now consider  $g' = \sum_{n=0}^{\infty} d_n \tau^n \in R\{\{\tau\}\}$ , where  $d_0 = b_0^{-1}$  and we inductively define  $d_n = -b_0^{-1} (b_1 d_{n-1}^{q^1} + b_2 d_{n-2}^{q^2} + \cdots + b_n d_0^{q^n})$ . Then as above, one can easily check that  $f \circ g' = \mathbb{1}$  and hence it is a right inverse of  $f$  in  $R\{\{\tau\}\}$ . But using the

associativity property of  $R\{\{\tau\}\}$  we get  $g' = (g \circ f) \circ g' = g \circ (f \circ g') = g$  and hence  $g$  is both a left and right inverse of  $f$  in  $S^\dagger$ .  $\square$

**Proposition 5.4.** *Let  $B$  denote the subring  $\mathbb{F}_q[t] \subseteq A$ . Let  $f : E \rightarrow G$  be a  $B$ -linear homomorphism of admissible  $A$ -module schemes over  $\mathrm{Spf} R$ . Then  $f$  is  $A$ -linear.*

*Proof.* Given any element  $a \in A$ , we will show  $\varphi_G(a) \circ f = f \circ \varphi_E(a)$ . Both sides are  $B$ -linear homomorphisms  $E \rightarrow G$ ; indeed,  $f$  is  $B$ -linear by assumption, and both  $\varphi_G(a)$  and  $\varphi_E(a)$  are  $B$ -linear because  $A$  is commutative. Furthermore, on tangent spaces,  $\varphi_G(a) \circ f$  is multiplication by  $af'(0)$ , and  $f \circ \varphi_E(a)$  is multiplication by  $f'(0)a$ ; this is because the  $A$ -module schemes are admissible. Thus the two morphisms agree on tangent spaces and therefore they agree, by proposition 5.1.  $\square$

In other words, the forgetful functor from admissible  $A$ -modules schemes over  $R$  to admissible  $B$ -module schemes over  $R$  is fully faithful. This remains true if we allow  $B$  to be not just  $\mathbb{F}_q[t]$  but any sub- $\mathbb{F}_q$ -algebra of  $A$  strictly containing  $\mathbb{F}_q$ .

**Lemma 5.5.** *If  $q \geq 3$ , then  $q^i - q^{i-j} - j - 1 \geq 0$  for all  $j = 1, \dots, i$ .*

*Proof.* Consider  $f(x) = q^i - q^{i-x} - x - 1$ , for  $1 \leq x \leq i$ . Then  $f(1) \geq 0$  since  $q \geq 3$ . Now  $f'(x) = q^{i-x} \ln q - 1$ . Since  $\ln q > 1$  for  $q \geq 3$ , we have  $f'(x) \geq 0$  for all  $1 \leq x \leq i$  and hence  $f(x) \geq 0$  for all  $1 \leq x \leq i$  and we are done.  $\square$

**Theorem 5.6.** *Suppose  $q \geq 3$  and  $v(a_i) \geq q^i - 1$ , for all  $i \geq 1$ . Then there exists a unique  $A$ -linear homomorphism  $f : E \rightarrow \hat{\mathbb{G}}_a$ , written  $f = \sum_{i=0}^{\infty} b_i \tau^i$  in coordinates, such that  $v(b_i) \geq i$  and  $b_0 = 1$ . Moreover,  $f$  is an isomorphism of  $A$ -module schemes over  $S$ .*

*Proof.* Let  $f = \sum_{i=0}^{\infty} b_i \tau^i$ ,  $b_i \in R$ , where  $b_0 = 1$  and

$$(5.3) \quad b_i = \pi^{-1} (1 - \pi^{q^i - 1})^{-1} \sum_{j=1}^i b_{i-j} a_j^{q^{i-j}}.$$

Indeed, this is the only possible choice for  $f$ , by proposition 5.2. Conversely, it is easy to see that  $f$  satisfies  $\varphi(t) \circ f = f \circ \varphi(t)$ , which implies  $\varphi(b) \circ f = f \circ \varphi(b)$  for all  $b \in B$ .

Let us now show  $v(b_i) \geq i$ . For  $i = 0$ , it is clear. For  $i \geq 1$ , we may assume by induction that  $v(b_j) \geq j$  for all  $j = 1, \dots, i-1$ . By (5.3), we have  $v(b_i) \geq \min\{v(b_{i-j} a_j^{q^{i-j}}) - 1 \mid j = 1, \dots, i\}$ . Now

$$\begin{aligned} v(b_{i-j} a_j^{q^{i-j}}) - 1 &\geq v(b_{i-j}) + v(a_j^{q^{i-j}}) - 1 \\ &\geq i - j + q^{i-j} (q^j - 1) - 1 \\ &= i - j + q^i - q^{i-j} - 1 \\ &\geq i, \text{ by lemma 5.5.} \end{aligned}$$

Therefore we have  $v(b_i) \geq i$ .

Therefore  $f$  is a restricted power series and hence defines a map between  $\pi$ -formal schemes  $f : E \rightarrow \hat{\mathbb{G}}_a$  which is  $A$ -linear.

Let us show that  $f$  is an isomorphism. By proposition 5.3, there exists a linear map  $g : \hat{\mathbb{G}}_a \rightarrow E$  such that  $f \circ g = g \circ f = \mathbb{1}$ . Then  $g$  is also  $A$ -linear for formal

reasons. Indeed, for any  $a \in A$ , we have  $f(g(\varphi(a)x)) = \varphi(a)x = f(\varphi(a)g(x))$ . Since  $f$  is injective, we must have  $g(\varphi(a)x) = \varphi(a)g(x)$  which shows the  $A$ -linearity  $g$  and we are done.  $\square$

We conclude this section with a useful observation which will not however be used here. Letting  $\mathbb{G}_a^{\text{for}}$  denote the formal completion of  $\hat{\mathbb{G}}_a$  along the identity section  $\text{Spf } R \rightarrow \hat{\mathbb{G}}_a$ . Thus we have  $\mathbb{G}_a^{\text{for}} = \text{Spf } R[[x]]$ , where  $R[[x]]$  has the  $(\pi, x)$ -adic topology. We want to extend the  $A$ -action on  $\mathbb{G}_a^{\text{for}}$  to an action of  $\hat{A}$ :

$$(5.4) \quad \hat{A} \rightarrow \text{End}_{\mathbb{F}_q}(\mathbb{G}_a^{\text{for}}/S).$$

Recall that  $\text{End}_{\mathbb{F}_q}(\mathbb{G}_a^{\text{for}})$  agrees with the non-commutative power-series ring  $R\{\{\tau\}\}$ , with commutation law  $\tau b = b^q \tau$  for  $b \in R$ . Therefore for any  $a \in A$ , we can write

$$\varphi(a) = \sum_j \alpha_j \tau^j$$

where  $\alpha_j \in R$ . Each  $\alpha_j$  can be thought of as a function of  $a \in A$ . To construct (5.4) it is enough to prove that these functions are  $\mathfrak{p}$ -adically continuous, which also implies that such an extension to a continuous  $\hat{A}$ -action is unique. This is a consequence of the following result.

**Proposition 5.7.** *If  $a \in \mathfrak{p}^n$ , then  $\alpha_j \in \mathfrak{p}^{n-j}R$ .*

*Proof.* Clearly, it is true for  $n = 0$ . Now assume it is true for some given  $n$ . Suppose  $a \in \mathfrak{p}^{n+1}$  and write  $a = \pi b$ , where  $b \in \mathfrak{p}^n$ . Let  $\varphi(b) = \sum_j \beta_j \tau^j$  and  $\varphi(\pi) = \sum_k \gamma_k \tau^k$ . Then we have

$$\sum_j \alpha_j \tau^j = \varphi(a) = \varphi(\pi)\varphi(b) = \sum_k \gamma_k \tau^k \sum_j \beta_j \tau^j = \sum_{k,j} \gamma_k \beta_j^{q^k} \tau^{j+k}$$

and hence  $\alpha_j = \sum_{k=0}^j \gamma_k \beta_{j-k}^{q^k}$ . So to show  $\alpha_j \in \mathfrak{p}^{n+1-j}R$ , it suffices to show

$$\gamma_k \beta_{j-k}^{q^k} \in \mathfrak{p}^{n+1-j}R, \text{ for } 0 \leq k \leq j \leq n+1.$$

By induction we have  $\beta_{j-k} \in \mathfrak{p}^{n-(j-k)}R$  and hence  $\gamma_k \beta_{j-k}^{q^k} \in \mathfrak{p}^{(n-(j-k))q^k}R$ . Since we have  $(n-(j-k))q^k \geq n-j+1$  for  $k \geq 1$ , we then have  $\gamma_k \beta_{j-k}^{q^k} \in \mathfrak{p}^{n-j+1}R$ . For  $k = 0$ , because  $\varphi$  is a strict module structure, we have  $\gamma_0 = \pi$  and hence  $\gamma_0 \beta_j \in \pi \mathfrak{p}^{n-j}R = \mathfrak{p}^{1+n-j}R$ .  $\square$

## 6. CHARACTERS OF $N^n$ —UPPER BOUNDS

We continue to let  $E$  denote the admissible  $A$ -module scheme over  $S$  of (5.1).

**Lemma 6.1.** *For all  $n \geq 0$ ,  $\phi^n(x) = \pi^n x^{(n)} + O(n-1)$ , where  $O(n-1)$  are elements of order less than equal to  $n-1$ .*

*Proof.* For  $n = 0$ , it is clear. For  $n \geq 1$ , we have by induction

$$\begin{aligned} \phi^n(x) &= \phi(\pi^{n-1}x^{(n-1)} + O(n-2)) \\ &= \pi^{n-1}\phi(x^{(n-1)}) + O(n-1) \\ &= \pi^{n-1}\pi\delta(x^{(n)}) + (x^{(n-1)})^{\hat{q}} + O(n-1) \\ &= \pi^n x^{(n)} + O(n-1). \end{aligned}$$

□

**Theorem 6.2.** *Assume  $q \geq 3$ . For any  $n \geq 0$ , let  $H^n$  denote the kernel of the projection  $u : J^{n+1}E \rightarrow J^n E$ . Then there is a unique  $A$ -linear isomorphism  $\vartheta_n : H^n \rightarrow \hat{\mathbb{G}}_a$  of the form  $\vartheta_n(x) = x + b_1 x^q + b_2 x^{q^2} + \dots$  in coordinates such that  $v(b_i) \geq i$  for all  $i \geq 1$ . Further, the  $R$ -module  $\text{Hom}_A(H^n, \hat{\mathbb{G}}_a)$  is freely generated by  $\vartheta_n$*

*Proof.* First observe that we have

$$\begin{aligned} \varphi_E(t)\phi^n(x) &= \phi^n(\varphi_E(t)) \\ &= \phi^n(\pi)\phi^n(x) + \phi^n(a_1)\phi^n(x)^q + \dots + \phi^n(a_r)\phi^n(x)^{q^r}. \end{aligned}$$

Second, the subscheme  $H^n$  is defined by setting the  $x, x', \dots, x^{(n-1)}$  coordinates to 0. Combining these two observations and lemma 6.1, we obtain

$$\pi^n \varphi_E(t)x^{(n)} = \pi \pi^n x^{(n)} + \phi^n(a_1)(\pi^n x^{(n)})^q + \dots + \phi^n(a_r)(\pi^n x^{(n)})^{q^r}$$

and hence

$$\varphi_E(t)x^{(n)} = \pi x^{(n)} + \phi^n(a_1)\pi^{n(q-1)}(x^{(n)})^q + \dots + \phi^n(a_r)\pi^{n(q^r-1)}(x^{(n)})^{q^r}.$$

But then by theorem 5.6, there is a unique isomorphism  $(H^n, \varphi_{H^n}) \rightarrow (\hat{\mathbb{G}}_a, \varphi_{\hat{\mathbb{G}}_a})$  of the kind desired.

Finally it follows from corollary 5.2 that  $\text{Hom}_A(H^n, \hat{\mathbb{G}}_a)$  is freely generated as an  $R$ -module by  $\vartheta_n$ . □

We emphasize that for  $q = 2$ , it is not generally true that  $H^n$  is isomorphic to  $\hat{\mathbb{G}}_a$ . We do however have an argument along similar lines that  $\text{Hom}_A(H^n, \hat{\mathbb{G}}_a)$  is free of rank 1 over  $R$ , but to keep the exposition short, we have not included it here.

Now consider the exact sequence

$$0 \rightarrow H^n \rightarrow N^n \rightarrow N^{n-1} \rightarrow 0$$

and the corresponding long exact sequence

$$0 \rightarrow \text{Hom}_A(N^{n-1}, \hat{\mathbb{G}}_a) \rightarrow \text{Hom}_A(N^n, \hat{\mathbb{G}}_a) \rightarrow \text{Hom}_A(H^n, \hat{\mathbb{G}}_a) \rightarrow \dots$$

By the theorem above, the image of the map  $\text{Hom}_A(N^n, \hat{\mathbb{G}}_a) \rightarrow \text{Hom}_A(H^n, \hat{\mathbb{G}}_a)$  can be regarded as a sub- $R$ -module of  $R$ . Therefore we have a filtration of  $R$ -modules

$$\text{Hom}_A(N^n, \hat{\mathbb{G}}_a) \supseteq \text{Hom}_A(N^{n-1}, \hat{\mathbb{G}}_a) \supseteq \dots \supseteq \text{Hom}_A(N^0, \hat{\mathbb{G}}_a) = 0,$$

and each associated graded module is canonically a submodule of  $R$ .

In particular, we have the following:

**Proposition 6.3.** *If  $R$  is a discrete valuation ring, then  $\text{Hom}_A(N^n, \hat{\mathbb{G}}_a)$  is a free  $R$ -module of rank at most  $n$ .*



7. THE LATERAL FROBENIUS AND CHARACTERS OF  $N^n$ 

Now we will construct a family of important operators which we call the *lateral Frobenius* operators. That is, for all  $n$ , we will construct maps  $\mathfrak{f} : N^{n+1} \rightarrow N^n$  which are lifts of Frobenius relative to the projections  $u : N^{n+1} \rightarrow N^n$  and hence make the system  $\{N^n\}_{n=0}^\infty$  into a prolongation sequence. Do note that *a priori* the  $A$ -modules  $N^n$  do not form a prolongation sequence to start with.

Let  $N^\infty$  denote the inverse limit the projection maps  $u : N^{n+1} \rightarrow N^n$ . Then the maps  $\mathfrak{f}$  induce a lift of Frobenius on  $N^\infty$ . Similarly on  $J^\infty E = \lim_n J^n E$ , the maps  $\phi$  induce a lift of Frobenius. Now for all  $n \geq 1$ , the inclusion  $N^n \hookrightarrow J^n E$  is a closed immersion and hence induces a closed immersion of schemes  $N^\infty \hookrightarrow J^\infty E$ . But  $\mathfrak{f}$  is not obtained by restricting  $\phi$  to  $N^\infty$ . In fact,  $\phi$  does not even preserve  $N^\infty$ . So  $\mathfrak{f}$  is an interesting operator which is distinct from  $\phi$ , although it does satisfy a certain relation with  $\phi$  which we will explain below.

Here we would also like to remark that the lateral Frobenius can also be constructed in the mixed-characteristic setting of  $p$ -jet spaces of arbitrary schemes. But it is much more involved than for Drinfeld modules, and the authors will present that theory in a subsequent note.

Let  $F : W_n \rightarrow W_{n-1}$  and  $V : W_{n-1} \rightarrow W_n$  denote the Frobenius and Verschiebung maps of 3.3. Let us arrange them in the following diagram, although it does not commute.

$$\begin{array}{ccc} W_n & \xrightarrow{V} & W_{n+1} \\ F \downarrow & & \downarrow F \\ W_{n-1} & \xrightarrow{V} & W_n \\ & & \downarrow F \\ & & W_{n-1} \end{array}$$

Rather the following is true

$$(7.1) \quad FFV = FVF.$$

Indeed, the operator  $FV$  is multiplication by  $\pi = \theta(t)$ , and  $F$  is a morphism of  $A$ -algebras.

We can re-express this in terms of jet spaces using the natural identifications  $J^n E \simeq W_n$  and  $N^n \simeq W_{n-1}$ . For jet spaces, let us switch to the notation  $i := V$ ,  $\phi := F$  for the right column, and  $\mathfrak{f} := F$  for the left column. Then the diagram above becomes the following:

$$\begin{array}{ccc} N^{n+1} & \xrightarrow{i} & J^{n+1} E \\ \mathfrak{f} \downarrow & & \downarrow \phi \\ N^n & \xrightarrow{i} & J^n E \\ & & \downarrow \phi \\ & & J^{n-1} E \end{array}$$

Note again that it is not commutative. However rewriting (7.1) in the above notation, we do have

$$(7.2) \quad \phi^{\circ 2} \circ i = \phi \circ i \circ \mathfrak{f}.$$

We emphasize that when we use the notation  $N^n$ , the  $A$ -module structure will always be understood to be the one that makes  $i$  an  $A$ -linear morphism. It should not be confused with the  $A$ -module structure coming by transport of structure from the isomorphism  $N^n \simeq W_{n-1} = J^{n-1}E$  of group schemes.

We also emphasize that while  $i$  is a morphism of  $S$ -schemes, the vertical arrows  $\phi$  and  $\mathfrak{f}$  in the diagram above lie over the Frobenius endomorphism  $\phi$  of  $S$ , rather than the identity morphism.

**Lemma 7.1.** *For any torsion-free  $R$ -algebra  $B$ , the map  $FV : W_n(B) \rightarrow W_n(B)$  is injective.*

*Proof.* Since  $B$  is torsion free, the ghost map  $W_n(B) \rightarrow B \times \cdots \times B$  is injective, and hence  $W_n(B)$  is torsion free. The result then follows because  $FV$  is multiplication by  $\pi$ .  $\square$

**Proposition 7.2.** *The morphism  $\mathfrak{f} : N^n \rightarrow N^{n-1}$  is  $A$ -linear.*

*Proof.* Since both  $\phi$  and  $i$  are  $A$ -linear morphisms, so are  $\phi i$  and  $\phi^2 i$ . Therefore for all  $a \in A$ , we have

$$\phi i(\mathfrak{f}(ax)) = \phi^2 i(ax) = a\phi^2 i(x) = a\phi i(\mathfrak{f}(x)) = \phi i(a\mathfrak{f}(x))$$

Thus the two morphisms  $N^{n+1} \rightarrow N^n$ , given by  $x \mapsto a\mathfrak{f}(x)$  and by  $x \mapsto \mathfrak{f}(ax)$ , become equal after application of  $\phi i$ . We can interpret the morphisms as two elements of  $N^n(B)$ , where  $B$  is the algebra representing the functor  $N^{n+1}$ , which become equal after applying  $\phi i$ . But since  $B$  is torsion free, lemma 7.1 implies these two elements must be equal.  $\square$

For  $0 \leq i \leq k-1$ , let us abusively write  $\mathfrak{f}^{\circ i}$  for the following composition

$$\mathfrak{f}^{\circ i} : N^n \xrightarrow{\overbrace{\mathfrak{f} \circ \cdots \circ \mathfrak{f}}^{i\text{-times}}} N^{n-i} \xrightarrow{u} N^{n-k}.$$

Then for all  $1 \leq i \leq n$ , we define  $\Psi_i \in \text{Hom}_A(N^n, \hat{\mathbb{G}}_a)$  as

$$(7.3) \quad \Psi_i = \vartheta_1 \circ \mathfrak{f}^{\circ i-1}$$

where  $\vartheta_1$  is as in theorem 6.2. Clearly, the maps  $\Psi_i$  are  $A$ -linear since each one of the maps above is. Finally, given a character  $\Psi \in \text{Hom}_A(N^{n-1}, \hat{\mathbb{G}}_a)$ , we will write  $\mathfrak{f}^* \Psi = \Psi \circ \mathfrak{f}$ .

The points of  $J^n E$  contained in  $N^n$  are those with Witt coordinates of the form  $(0, x_1, x_2, \dots, x_n)$ . We will use the abbreviated coordinates  $(x_1, \dots, x_n)$  on  $N^n$  instead.

**Lemma 7.3.** *For  $q \geq 3$  and all  $i = 1, \dots, n$ , we have*

$$\Psi_i(x_1, \dots, x_n) \equiv x_1^{\hat{q}^{i-1}} \pmod{\pi}.$$

*Proof.* Since  $\mathfrak{f}$  is identified with the Frobenius map  $F : W_n \rightarrow W_{n-1}$ , it reduces modulo  $\pi$  to the  $\hat{q}$ -th power of the projection map. Therefore, we have

$$\Psi_i(x_1, \dots, x_n) = \vartheta_1 \circ \mathfrak{f}^{\circ(i-1)}(x_1, \dots, x_n) \equiv \vartheta_1(x_1^{\hat{q}^{i-1}}) \pmod{\pi},$$

and hence is equivalent to  $x_1^{\hat{q}^{i-1}}$  modulo  $\pi$ , by the defining property of  $\vartheta_1$  in theorem 6.2.  $\square$

**Proposition 7.4.** *If  $R$  is a discrete valuation ring, then the elements  $\Psi_1, \dots, \Psi_n$  form an  $R$ -basis for  $\text{Hom}_A(N^n, \hat{\mathbb{G}}_a)$ .*

*Proof.* (For  $q \geq 3$ .) By proposition 6.3, the  $R$ -module  $\text{Hom}_A(N^n, \hat{\mathbb{G}}_a)$  is free of rank at most  $n$ . So to show the elements  $\Psi_1, \dots, \Psi_n$  form a basis, it is enough by Nakayama's lemma to show they are linearly independent modulo  $\pi$ . But for  $q \geq 3$ , by lemma 7.3, we have  $\Psi_i \equiv x_1^{\hat{q}^{i-1}} \pmod{\pi}$ , and so the  $\Psi_i$  map to linearly independent elements of  $R/\pi R \otimes_R \mathcal{O}(N^n)$ . Thus they are linearly independent in  $R/\pi R \otimes_R \text{Hom}_A(N^n, \hat{\mathbb{G}}_a)$ .  $\square$

## 8. $\mathbf{X}_\infty(E)$

We now assume further that  $R$  is a discrete valuation ring and  $E$  is a Drinfeld module over  $\text{Spf } R$ . Let  $r$  denote the rank of  $E$ . We continue to write  $\varphi_E(t) = a_0\tau^0 + a_1\tau^1 + \dots + a_r\tau^r$ , where  $a_0 = \pi$ ,  $a_i \in R$  for all  $i$ , and  $a_r \in R^*$ .

Given such a Drinfeld module, one of the important  $\delta$ -arithmetic objects that one can attach to it is the group of all  $\delta$ -characters of  $E$  to  $\hat{\mathbb{G}}_a$ , denoted  $\mathbf{X}_\infty(E)$ . In the case of elliptic curves, Buium has shown that this group contains important arithmetic data as analogues of Manin maps in differential algebra and has found diophantine applications on Heegner points on modular curves [BP2].

In this section and the next, we will determine the structure of  $\mathbf{X}_\infty(E)$ . In the case of elliptic curves, it falls in two distinct cases as to when the elliptic curve admits a canonical lift and when not. A similar story happens in our case when  $E$  is a Drinfeld module of rank 2, which one might consider the closest analogue of an elliptic curve. However, when the rank exceeds 2, the behavior of  $\mathbf{X}_\infty(E)$  offers much more interesting cases which leads us to introduce the concept of the *splitting order*  $m$  of a Drinfeld module  $E$ . This natural number is always less than or equal to the rank of  $E$  and when the rank equals 2, the notion coincides with the canonical lift property of Drinfeld modules.

We would like to point out here that our structure result for  $\mathbf{X}_\infty(E)$  is an integral version of that of [Bui2]. Buium shows that  $\mathbf{X}_\infty(E) \otimes_R K$  is generated by a single element as a  $K\{\phi^*\}$ -module where  $K = R[\frac{1}{p}]$ . But here we show that the module  $\mathbf{X}_\infty(E)$  itself is generated by a single element as a  $R\{\phi^*\}$ -module. Although our result is for Drinfeld modules over function rings in positive characteristic, our methods work in the elliptic curves over  $p$ -adic rings setting and hence this stronger result can be achieved in that case too.

The following theorem should be viewed as an analogue of the fact that an elliptic curve has no non-zero homomorphism of  $\mathbb{Z}$ -module schemes to  $\mathbb{G}_a$ . In our case, we show that no Drinfeld module admits a non-zero homomorphism of  $A$ -module schemes to  $\hat{\mathbb{G}}_a$ .

**Theorem 8.1.** *We have  $\mathbf{X}_0(E) = \{0\}$ .*

*Proof.* Any character  $f = \sum_{i \geq 0} b_i \tau^i \in \mathbf{X}_0(E)$  satisfies the following chain of equalities:

$$\begin{aligned} \varphi_{\hat{\mathbb{G}}_a}(t) \circ f &= f \circ \varphi_E(t) \\ \theta(t) \tau^0 \circ \sum_{i \geq 0} b_i \tau^i &= \sum_{i \geq 0} b_i \tau^i \circ \sum_j a_j \tau^j \\ \sum_{i \geq 0} \theta(t) b_i \tau^i &= \sum_{i \geq 0} \left( \sum_{j=0}^r b_{i-j} a_j^{q^{i-j}} \right) \tau^i \end{aligned}$$

Comparing the coefficients of  $\tau^i$  for  $i > r$ , we have

$$(8.1) \quad b_i (1 - \theta(t)^{q^{i-1}}) \theta(t) = a_r^{q^{i-r}} b_{i-r} + a_{r-1}^{q^{i-r+1}} b_{i-r+1} + \cdots + a_1^{q^{i-1}} b_{i-1}$$

Suppose  $f$  is nonzero. Then there exists an  $N$  such that  $b_{N-r} \neq 0$  and  $v(b_{N-r}) < v(b_i)$  for all  $i \geq N - r + 1$ . Then the valuation of the right-hand side of equation (8.1) for  $i = N$  becomes  $v(a_r^{q^{i-r}} b_{N-r}) = v(b_{N-r})$ , since  $v(a_r) = 0$ . But then taking the valuation of both sides of (8.1), we have

$$v(b_N) = v(b_{N-r}) - 1 < v(b_{N-r})$$

and  $N \geq N - r + 1$ , which is a contradiction to the hypothesis above. Therefore  $f$  must be 0.  $\square$

As a consequence the short exact sequence of  $A$ -module schemes over  $S$

$$(8.2) \quad 0 \rightarrow N^n \xrightarrow{i} J^n E \rightarrow E \rightarrow 0,$$

induces an exact sequence

$$(8.3) \quad 0 \rightarrow \mathbf{X}_n(E) \xrightarrow{i^*} \mathrm{Hom}_A(N^n, \hat{\mathbb{G}}_a) \xrightarrow{\partial} \mathrm{Ext}_A(E, \hat{\mathbb{G}}_a),$$

which we will use repeatedly.

The following result is the analogue of Buium's in the mixed-characteristic setting.

**Theorem 8.2.** *Let  $(E, \varphi_E)$  be a Drinfeld module of rank  $r$ .*

- (1)  $\mathbf{X}_r(E)$  is nonzero.
- (2) We have

$$\mathbf{X}_1(E) \simeq \begin{cases} R, & \text{if } E \text{ has a lift of Frobenius,} \\ \{0\}, & \text{otherwise.} \end{cases}$$

*Proof.* (1): Consider the exact sequence (8.3). By proposition 7.4, the  $R$ -module  $\mathrm{Hom}_A(N^n, \hat{\mathbb{G}}_a)$  is free of rank  $n$ . But also  $\mathrm{Ext}_A(E, \hat{\mathbb{G}}_a)$  is free of rank  $r - 1$ , by [Ge1]. Therefore when  $n = r$ , the kernel  $\mathbf{X}_n(E)$  is nonzero.

(2) Now consider  $\mathbf{X}_1(E)$ . It is contained in  $\mathrm{Hom}_A(N^1, \hat{\mathbb{G}}_a)$ , which is free of rank 1, and the quotient is contained in  $\mathrm{Ext}_A(E, \hat{\mathbb{G}}_a)$ , which is torsion free. Therefore  $\mathbf{X}_1(E)$  is either  $\{0\}$  or all of  $\mathrm{Hom}_A(N^1, \hat{\mathbb{G}}_a) \simeq R$ .

Let  $\mathbb{1}$  denote the identity map in  $\text{Hom}_A(\hat{\mathbb{G}}_a, \hat{\mathbb{G}}_a)$ . Then its image  $\partial(\mathbb{1})$  in  $\text{Ext}_A(E, \hat{\mathbb{G}}_a)$  is the class of the extension (8.2). Therefore we have the equivalences  $\mathbf{X}_1(E) \simeq R \iff i^*$  is an isomorphism  $\iff \partial(\mathbb{1}) = 0 \iff$  (8.2) is split  $\iff E$  has a lift of Frobenius.  $\square$

Define the *splitting order* of the Drinfeld module  $E$  to be the integer  $m$  such that  $\mathbf{X}_m(E) \neq \{0\}$  and  $\mathbf{X}_{m-1}(E) = \{0\}$ . We also say that  $E$  *splits at order*  $m$ . By the theorems above, we have  $1 \leq m \leq r$  and additionally  $m = 1$  if and only if  $E$  is a canonical lift.

**8.1. Splitting of  $J^n(E)$ .** The exact sequence (8.3) is split by the Teichmüller section  $v : E \rightarrow J^n E$ , as defined in section 3. We emphasize that  $v$  is only a morphism of  $\mathbb{F}_q$ -module schemes and is not a morphism of  $A$ -module schemes. Nevertheless, it induces an isomorphism

$$J^n(E) \xrightarrow{\sim} E \times N^n$$

of  $\mathbb{F}_q$ -module schemes. Therefore for any character  $\Theta \in \mathbf{X}_n(E)$ , we can write  $\Theta = g \oplus \Psi$  or

$$(8.4) \quad \Theta(x_0, \dots, x_n) = g(x_0) + \Psi(x_1, \dots, x_n),$$

where  $\Psi = i^* \Theta \in \text{Hom}_A(N^n, \hat{\mathbb{G}}_a)$  and  $g = v^* \Theta$ . Note that because  $v$  is only  $\mathbb{F}_q$ -linear,  $g$  is also only  $\mathbb{F}_q$ -linear. But it can still be expressed as an additive restricted power series.

**Lemma 8.3.** *For any  $R$ -algebra  $B$ , consider the exact sequence for all  $n \geq 1$*

$$0 \rightarrow W_{n-1}(B) \xrightarrow{V} W_n(B) \xrightarrow{T^n} B \rightarrow 0$$

*Then there exists a map  $g : B \rightarrow W_n(B)$  such that*

$$\begin{array}{ccc} W_n(B) & \xrightarrow{T^n} & B \\ \downarrow FV - VF & \swarrow g & \\ W_n(B) & & \end{array}$$

*commutes. It is of the form  $g(x) = (\pi x, c_1 x^{\hat{q}}, c_2 x^{\hat{q}^2}, \dots)$ , for some elements  $c_j \in R$ .*

*Proof.* For any  $y \in W_{n-1}(B)$ , we have

$$(FV - VF)(Vy) = FVVy - VVfy = \pi Vy - V(\pi y) = 0.$$

So such a function  $g$  exists.

To conclude that  $g(x)$  is of the given form, we use a homogeneity argument. Let  $(z_0, z_1, \dots)$  denote the ghost components of  $(x_0, x_1, \dots)$ . If interpret each  $x_j$  as an indeterminate of degree  $\hat{q}^j$ , then each  $z_j$  is a homogenous polynomial in the  $x_0, \dots, x_j$  of degree  $\hat{q}^j$  and with coefficients in  $A$ :  $z_1 = x_0^{\hat{q}} + \pi x_1$ , and so on. Solving for  $x_j$  in terms of  $z_0, \dots, z_j$ , we see that  $x_j$  is a homogenous polynomial in the  $z_0, \dots, z_j$  with coefficients in  $A[1/\pi]$ .

Now let  $(y_0, y_1, \dots)$  denote  $(FV - VF)(x_0, x_1, \dots)$ , where  $y_j \in R[x_0, \dots, x_j]$ . Then the ghost components of  $(y_0, y_1, \dots)$  are  $(\pi z_0, 0, 0, \dots) = (\pi x_0, 0, 0, \dots)$ . It follows that  $y_0 = \pi x_0$ . Further, by the above,  $y_j$  is an element of  $R[x_0, \dots, x_j]$  but

also a homogeneous polynomial in  $\pi x_0$  of degree  $\hat{q}^j$  and with coefficients in  $A[1/\pi]$ . Therefore it is of the form  $c_j x_0^{\hat{q}^j}$  for some  $c_j \in R$ .  $\square$

**Proposition 8.4.** *Let  $\Theta$  be a character in  $\mathbf{X}_n(E)$ .*

(1) *We have*

$$i^* \phi^* \Theta = \mathfrak{f}^*(i^* \Theta) + \gamma \Psi_1,$$

where  $\gamma = \pi g'(0)$  and  $g'(x_0)$  denotes the usual derivative of the polynomial  $g(x_0) \in R[x_0]$  of equation (8.4).

(2) *For  $n \geq 1$ , we have*

$$i^*(\phi^{\circ n})^* \Theta = (\mathfrak{f}^{n-1})^* i^* \phi^* \Theta.$$

*Proof.* (1): By lemma 8.3, we have

$$(\phi \circ i - i \circ \mathfrak{f})(x_1, \dots, x_{n+1}) = (\pi x_1, c_1 x_1^{\hat{q}}, c_2 x_1^{\hat{q}^2}, \dots),$$

where  $c_j \in R$ . Therefore we have

$$(8.5) \quad \begin{aligned} ((i^* \phi^* - \mathfrak{f}^* i^*) \Theta)(x_1, \dots, x_{n+1}) &= \Theta(\pi x_1, c_1 x_1^{\hat{q}}, \dots) \\ &= g(\pi x_1) + \Psi(c_1 x_1^{\hat{q}}, \dots), \end{aligned}$$

where  $g$  and  $\Psi$  are as in equation (8.4). In particular, the character  $(i^* \phi^* - \mathfrak{f}^* i^*) \Theta$  depends only on  $x_1$ . Therefore it is of the form  $\gamma \Psi_1$ , for some  $\gamma \in R$ . Further since by theorem 6.2 we have  $\Psi_1'(0) = 1$ , the coefficient  $\gamma$  is simply the linear coefficient of  $(i^* \phi^* - \mathfrak{f}^* i^*) \Theta$ , which by (8.5) is  $\pi g'(0)$ .

(2): This is another way of expressing  $\phi^{\circ n} \circ i = \phi \circ i \circ \mathfrak{f}^{\circ(n-1)}$ , which follows from (7.2) by induction.  $\square$

**8.2. Frobenius and the filtration by order.** We would like to fix a notational convention here. Let  $u : J^n E \rightarrow J^{n'} E$  denote the canonical projection map for any  $n' < n$ , given in Witt coordinates by  $u(x_0, \dots, x_n) = (x_0, \dots, x_{n'})$ .

Consider the following morphism of exact sequences of  $A$ -modules

$$\begin{array}{ccccccccc} 0 & \longrightarrow & N^n & \xrightarrow{i} & J^n E & \xrightarrow{u} & E & \longrightarrow & 0 \\ & & \downarrow u & & \downarrow u & & \parallel & & \\ 0 & \longrightarrow & N^{n-1} & \xrightarrow{i} & J^{n-1} E & \xrightarrow{u} & E & \longrightarrow & 0 \end{array}$$

Since  $\mathbf{X}_0(E) = \{0\}$  by theorem 8.1, applying  $\text{Hom}_A(-, \hat{\mathbb{G}}_a)$  to the above, we obtain the following morphism of exact sequences of  $R$ -modules

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathbf{X}_n(E) & \xrightarrow{i^*} & \text{Hom}_A(N^n, \hat{\mathbb{G}}_a) & \xrightarrow{\partial} & \text{Ext}_A(E, \hat{\mathbb{G}}_a) \\ & & \uparrow u^* & & \uparrow u^* & & \parallel \\ 0 & \longrightarrow & \mathbf{X}_{n-1}(E) & \xrightarrow{i^*} & \text{Hom}_A(N^{n-1}, \hat{\mathbb{G}}_a) & \xrightarrow{\partial} & \text{Ext}_A(E, \hat{\mathbb{G}}_a) \end{array}$$

**Proposition 8.5.** *For any  $n \geq 0$ , the diagram*

$$\begin{array}{ccc} \mathbf{X}_n(E)/\mathbf{X}_{n-1}(E) & \xrightarrow{\phi^*} & \mathbf{X}_{n+1}(E)/\mathbf{X}_n(E) \\ \downarrow i^* & & \downarrow i^* \\ \mathrm{Hom}_A(N^n, \hat{\mathbb{G}}_a)/\mathrm{Hom}_A(N^{n-1}, \hat{\mathbb{G}}_a) & \xrightarrow[\sim]{\mathfrak{f}^*} & \mathrm{Hom}_A(N^{n+1}, \hat{\mathbb{G}}_a)/\mathrm{Hom}_A(N^n, \hat{\mathbb{G}}_a) \end{array}$$

*is commutative. The morphisms  $i^*$  and  $\phi^*$  are injective, and  $\mathfrak{f}^*$  is bijective.*

In fact, we will show in corollary 9.9 that all the morphisms in the diagram of proposition 8.5 are isomorphisms.

*Proof.* For  $n \geq 1$ , commutativity of the diagram follows from proposition 8.4; for  $n = 0$ , it follows from theorem 8.1.

The maps  $i^*$  are injective because the projections  $J^n E \rightarrow J^{n-1} E$  and  $N^n \rightarrow N^{n-1}$  have the same kernel, and  $\mathfrak{f}^*$  is an isomorphism by proposition 7.4. It follows that  $\phi^*$  is an injection.  $\square$

**8.3. The character  $\Theta_m$ .** Recall the exact sequence (8.3)

$$0 \rightarrow \mathbf{X}_n(E) \xrightarrow{i^*} \mathrm{Hom}_A(N^n, \hat{\mathbb{G}}_a) \xrightarrow{\partial} \mathrm{Ext}_A(E, \hat{\mathbb{G}}_a)$$

Let  $m$  denote the splitting order of  $E$ . Then for all  $n < m$ , the map

$$\partial : \mathrm{Hom}_A(N^n, \hat{\mathbb{G}}_a) \rightarrow \mathrm{Ext}_A(E, \hat{\mathbb{G}}_a)$$

is injective since  $\mathbf{X}_n(E) = \{0\}$ . But at  $n = m$ , we have  $\mathbf{X}_m(E) \neq \{0\}$ , and so there is a nonzero character  $\Psi \in \mathrm{Hom}_A(N^m, \hat{\mathbb{G}}_a)$  in the kernel of  $\partial$ . Write  $\Psi = \tilde{\lambda}_m \Psi_m - \tilde{\lambda}_{m-1} \Psi_{m-1} - \cdots - \tilde{\lambda}_1 \Psi_1$ , where  $\tilde{\lambda}_i \in R$  for all  $i = 0, \dots, m-1$ . Then we necessarily have  $\tilde{\lambda}_m \neq 0$  since  $\mathbf{X}_{m-1} = \{0\}$ . Therefore we have

$$(8.6) \quad \partial \Psi_m = \lambda_{m-1} \partial \Psi_{m-1} + \cdots + \lambda_1 \partial \Psi_1 \in \mathrm{Ext}_A(E, \hat{\mathbb{G}}_a)_K$$

where  $\lambda_i = \tilde{\lambda}_i / \tilde{\lambda}_m$  for all  $i = 1, \dots, m-1$ . This implies that the character

$$\Psi_m - \lambda_1 \Psi_{m-1} - \cdots - \lambda_{m-1} \Psi_1$$

is in  $\ker(\partial)$  and hence by the main exact sequence (8.3), there exists a unique  $\Theta_m \in \mathbf{X}_m(E)_K$  such that

$$(8.7) \quad i^* \Theta_m = \Psi_m - \lambda_{m-1} \Psi_{m-1} - \cdots - \lambda_1 \Psi_1$$

It then follows immediately that  $\Theta_m$  is a  $K$ -linear basis for  $\mathbf{X}_m(E)_K$ , say by propositions 7.4 and 8.5. (We will show in corollary 9.9 that  $\Theta_m$  actually lies in the group  $\mathbf{X}_m(E)$  of integral characters, and is in fact an integral basis for it.)

**Proposition 8.6.** *Let  $m$  denote the splitting order of  $E$ . Then for any  $j \geq 0$ , the character  $i^*(\phi^*)^j \Theta_m$  agrees with  $\Psi_{m+j}$  modulo rational characters of lower order, and the elements  $\Theta_m, \phi^* \Theta_m, \dots, \phi^{n-m} \Theta_m$  are a basis of the  $K$ -vector space  $\mathbf{X}_n(E)_K$ .*

*Proof.* By 8.5, each element  $\phi^{i^*} \Theta_m$  lies in  $\mathbf{X}_{m+i}(E)$  but not in  $\mathbf{X}_{m+i-1}(E)$ . Therefore such elements are linearly independent. At the same time, by the diagram above, each  $\mathbf{X}_{m+i}(E)/\mathbf{X}_{m+i-1}(E)$  has rank at most 1. Thus the rank of  $\mathbf{X}_n(E)$  is at most  $n - m + 1$ , and so the elements in question form a spanning set.  $\square$

Do note that this result will be improved to an integral version in theorem 9.10.

## 9. EXT GROUPS AND DE RHAM COHOMOLOGY

We will prove theorem 1.1 in this section. We continue with the notation from the previous section. In particular,  $R$  is a discrete valuation ring.

We will briefly describe our strategy in the next few lines. Recall from (8.7) the equality

$$i^* \Theta_m = \Psi_m - \lambda_{m-1} \Psi_{m-1} - \cdots - \lambda_1 \Psi_1$$

where  $\lambda_j \in K$ . *A priori*, the elements  $\lambda_j$  need not belong to  $R$ , but we prove in theorem 9.8 that they actually do. This will imply that  $i^* \Theta_m$  lies in  $\text{Hom}_A(N^m, \hat{\mathbb{G}}_a)$  and  $\ker(\partial)$ , and hence by the exact sequence (8.3), we have  $\Theta_m \in \mathbf{X}_m(E)$ —that is, the character  $\Theta_m$  is integral. From there, it is an easy consequence that  $\mathbf{X}_n(E)$  is generated by  $\Theta_m, \dots, \Theta_m^{\phi_m^{n-m}}$  as an  $R$ -module.

So the key result to prove is theorem 9.8. But it will require some preparation before we can present the proof. For all  $n \geq 1$ , we will define maps from  $\text{Hom}_A(N^n, \hat{\mathbb{G}}_a)$  to  $\text{Ext}^\sharp(E, \hat{\mathbb{G}}_a)$  which is also interpreted as the de Rham cohomology from associated to the Drinfeld module  $E$ . These maps are obtained by push-outs of  $J^n E$  by  $\Psi \in \text{Hom}_A(N^n, \hat{\mathbb{G}}_a)$ . To give an idea, do note that, for every  $n \geq 1$ , there are canonical elements  $E_\Psi^* \in \text{Ext}_A(E, \hat{\mathbb{G}}_a)$  where the  $E_\Psi^*$  is a push-out of  $J^n E$  by  $\Psi$  as follows

$$\begin{array}{ccccccccc} 0 & \longrightarrow & N^n & \xrightarrow{i} & J^n E & \xrightarrow{u} & E & \longrightarrow & 0 \\ & & \Psi \downarrow & & g_\Psi \downarrow & & \parallel & & \\ 0 & \longrightarrow & \hat{\mathbb{G}}_a & \longrightarrow & E_\Psi^* & \longrightarrow & E & \longrightarrow & 0 \end{array}$$

as  $E_\Psi^* \in \text{Ext}_A(E, \hat{\mathbb{G}}_a)$ . It leads to a very interesting theory of  $\delta$ -modular forms over the moduli space of Drinfeld modules and will be studied in a subsequent paper. And similar to previous cases, the main principles carry over to the case of elliptic curves or abelian schemes as well.

Now we introduce the theory of extensions of  $A$ -module group schemes. Given an extension  $\eta_C \in \text{Ext}(G, T)$  and  $f : T \rightarrow T'$  where  $G, T$  and  $T'$  are  $A$ -modules and  $f$  is an  $A$ -linear map we have the following diagram of the push-forward extension  $f_* C$ .

$$\begin{array}{ccccccccc} 0 & \longrightarrow & T & \longrightarrow & C & \longrightarrow & G & \longrightarrow & 0 \\ & & f \downarrow & & \downarrow & & \parallel & & \\ 0 & \longrightarrow & T' & \longrightarrow & f_* C & \longrightarrow & G & \longrightarrow & 0 \end{array}$$

The class of  $f_* C$  is obtained as follows—the class of  $\eta_C$  is represented by a linear (not necessarily  $A$ -linear) function  $\eta_C : G \rightarrow T$ . Then  $\eta_{f_* C}$  is represented by the class  $\eta_{f_* C} = [f \circ \eta_C] \in \text{Ext}(E, T')$ . In terms of the action of  $t \in A$ ,  $\varphi_C(t)$  is given by  $\begin{pmatrix} \varphi_C(t) & 0 \\ \eta_C & \varphi_T(t) \end{pmatrix}$  where  $\eta_C : G \rightarrow T$ . Then  $\varphi_{f_* C}(t)$  is given by

$$(9.1) \quad \begin{pmatrix} \varphi_G(t) & 0 \\ f(\eta_C) & \varphi_{T'}(t) \end{pmatrix}$$



Now consider the exact sequence

$$(9.2) \quad 0 \rightarrow N^n \xrightarrow{i} J^n E \xrightarrow{u} E \rightarrow 0$$

Given a  $\Psi \in \text{Hom}_A(N^n, \hat{\mathbb{G}}_a)$  consider the push out

$$\begin{array}{ccccccc} 0 & \longrightarrow & N^n & \xrightarrow{i} & J^n E & \xrightarrow{u} & E \longrightarrow 0 \\ & & \Psi \downarrow & & g_\Psi \downarrow & & \parallel \\ 0 & \longrightarrow & \hat{\mathbb{G}}_a & \xrightarrow{i} & E_\Psi^* & \longrightarrow & E \longrightarrow 0 \end{array}$$

where  $E_\Psi^* = \frac{J^n E \times \hat{\mathbb{G}}_a}{\Gamma(N^n)}$  and  $\Gamma(N^n) = \{(i(z), -\Psi(z)) \mid z \in N^n\} \subset J^n E \times N^n$  and  $g_\Psi(x) = [x, 0] \in E_\Psi^*$ .

The Teichmüller section  $v : E \rightarrow J^n(E)$  is an  $\mathbb{F}_q$ -linear splitting of the sequence (9.2). The induced retraction

$$\rho = \mathbb{1} - v \circ u : J^n(E) \rightarrow N^n$$

is given in coordinates simply by  $\rho : (x_0, \dots, x_n) \mapsto (x_1, \dots, x_n)$ . Let us denote by  $s_{\text{Witt}}$  the morphism on Lie algebras induced by  $\rho$ . Thus we have the following split exact sequence of  $R$ -modules

$$0 \longrightarrow \text{Lie } N^n \xrightleftharpoons[s_{\text{Witt}}]{Di} \text{Lie } J^n E \xrightarrow{Du} \text{Lie}(E) \longrightarrow 0$$

Let  $s_\Psi$  denote the induced splitting of the push out extension

$$0 \longrightarrow \text{Lie } \hat{\mathbb{G}}_a \xrightleftharpoons[s_\Psi]{g_\Psi} \text{Lie}(E_\Psi^*) \longrightarrow \text{Lie}(E) \longrightarrow 0$$

It is given explicitly by  $\tilde{s}_\Psi : \text{Lie } J^n E \times \text{Lie } \hat{\mathbb{G}}_a \rightarrow \text{Lie } \hat{\mathbb{G}}_a$

$$\tilde{s}_\Psi(x, y) := D\Psi(s_{\text{Witt}}(x)) + y$$

and

$$s_\Psi : \text{Lie}(E_\Psi^*) = \frac{\text{Lie } J^n E \times \text{Lie } \hat{\mathbb{G}}_a}{\text{Lie } \Gamma(N^n)} \rightarrow \text{Lie } \hat{\mathbb{G}}_a$$

This induces the following morphism of exact sequences

$$(9.3) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \mathbf{X}_n(E) & \longrightarrow & \text{Hom}_A(N^n, \hat{\mathbb{G}}_a) & \longrightarrow & \text{Ext}(E, \hat{\mathbb{G}}_a) \\ & & \downarrow & & \Psi \mapsto (E_\Psi^*, s_\Psi) \downarrow & & \parallel \\ 0 & \longrightarrow & \text{Lie}(E)^* & \longrightarrow & \text{Ext}^\sharp(E, \hat{\mathbb{G}}_a) & \longrightarrow & \text{Ext}(E, \hat{\mathbb{G}}_a) \longrightarrow 0 \end{array}$$

**Proposition 9.1.** *Let  $\Theta$  be a character in  $\mathbf{X}_n(E)$ , and put  $\Psi = i^*\Theta \in \text{Hom}_A(N^n, \hat{\mathbb{G}}_a)$  and  $g = v^*\Theta : E \rightarrow \hat{\mathbb{G}}_a$ , as in equation (8.4).*

- (1) *The map  $\mathbf{X}_n(E) \rightarrow \text{Lie}(E)^*$  of (9.3) sends  $\Theta$  to  $-Dg$ .*
- (2) *Let  $\tilde{\Theta} = \phi^*\Theta$ , and put  $\tilde{\Psi} = i^*\tilde{\Theta}$  and  $\tilde{g} = v^*\tilde{\Theta}$ . Then we have  $\tilde{g}(x) = g(x^q)$  and  $\tilde{\Psi}(y) = \Psi(\rho(\phi(i(y)))) + g(\pi y_1)$ .*

*Proof.* (1): Let us recall in explicit terms how the map is given. For the split extension  $E \times \hat{\mathbb{G}}_a$ , the retractions  $\text{Lie}(E) \times \text{Lie } \hat{\mathbb{G}}_a = \text{Lie}(E \times \hat{\mathbb{G}}_a) \rightarrow \text{Lie } \hat{\mathbb{G}}_a$  are in bijection with maps  $\text{Lie}(E) \rightarrow \text{Lie } \hat{\mathbb{G}}_a$ , a retraction  $s$  corresponding to map

$x \mapsto s(x, 0)$ . Therefore to determine the image of  $\Theta$ , we need to identify  $E_\Psi^*$  with a split extension and then apply this map to  $s_\Psi$ .

A trivialization of the extension  $E_\Psi^*$  is given by the map

$$\frac{J^n E \times \hat{\mathbb{G}}_a}{\Gamma(N^n)} = E_\Psi^* \xrightarrow{\sim} E \times \hat{\mathbb{G}}_a$$

defined by  $[a, b] \mapsto (u(a), \Theta(a) + b)$ . The inverse isomorphism  $H$  is then given by the expression

$$H(x, y) = [v(x), y - \Theta(v(x))],$$

and so the composition  $E \rightarrow E \times \hat{\mathbb{G}}_a \rightarrow E_\Psi^* \rightarrow \hat{\mathbb{G}}_a$  is simply  $-\Theta \circ v = -g$ , which induces the map  $-Dg$  on the Lie algebras.

(2): We have

$$\begin{aligned} \tilde{\Theta}(x) &= \Theta(\phi(x)) \\ &= \Psi(\rho(\phi(x))) + g(x_0^{\hat{q}} + \pi x_1) \\ &= (\Psi(\rho(\phi(x))) + g(\pi x_1)) + g(x_0^{\hat{q}}). \end{aligned}$$

In other words, we have  $\tilde{\Psi}(\rho(x)) = \Psi(\rho(\phi(x))) + g(\pi x_1)$  and  $\tilde{g}(x_0) = g(x_0^{\hat{q}})$ . Setting  $x = i(y)$ , we obtain the desired result.  $\square$

**Proposition 9.2.** *If  $\Psi \in i^* \phi^*(\mathbf{X}_n(E))$ , then the class  $(E_\Psi^*, s_\Psi) \in \text{Ext}^\sharp(E, \hat{\mathbb{G}}_a)$  is zero.*

*Proof.* We know from diagram (9.3) that  $E_\Psi^*$  is a trivial extension since  $\tilde{\Psi}$  lies in  $i^* \mathbf{X}_{n+1}(E)$ . Now as in part (2) of proposition 9.1, we have, in the notation of that proposition,  $\tilde{g}(x_0) = g(x_0^{\hat{q}})$  and hence  $D\tilde{g} = 0$ . Therefore by part (1) of that proposition, the class in  $\text{Ext}^\sharp(E, \hat{\mathbb{G}}_a)$  is zero.  $\square$

**9.1. The crystal  $\mathbf{H}(E)$ .** The  $\phi$ -linear map  $\phi^* : \mathbf{X}_{n-1}(E) \rightarrow \mathbf{X}_n(E)$  induces a linear map  $\mathbf{X}_{n-1}(E)^\phi \rightarrow \mathbf{X}_n(E)$ , which we will abusively also denote  $\phi^*$ . We then define

$$\mathbf{H}_n(E) = \frac{\text{Hom}_A(N^n, \hat{\mathbb{G}}_a)}{i^* \phi^*(\mathbf{X}_{n-1}(E)^\phi)}$$

Then  $u : N^{n+1} \rightarrow N^n$  induces  $u^* : \text{Hom}_A(N^n, \hat{\mathbb{G}}_a) \rightarrow \text{Hom}_A(N^{n+1}, \hat{\mathbb{G}}_a)$ . And since  $u^* i^* \phi^*(\mathbf{X}_n(E)) = i^* u^* \phi^*(\mathbf{X}_n(E)) = i^* \phi^* u^*(\mathbf{X}_n(E)) \subset i^* \phi^*(\mathbf{X}_{n+1}(E))$ , it also induces a map  $u^* : \mathbf{H}_n(E) \rightarrow \mathbf{H}_{n+1}(E)$ . Define  $\mathbf{H}(E) = \lim_{\rightarrow} \mathbf{H}_n(E)$ .

Similarly,  $\mathfrak{f} : N^{n+1} \rightarrow N^n$  induces  $\mathfrak{f}^* : \text{Hom}_A(N^n, \hat{\mathbb{G}}_a) \rightarrow \text{Hom}_A(N^{n+1}, \hat{\mathbb{G}}_a)$ , which descends to a  $\phi$ -linear morphism of  $R$ -modules

$$\mathfrak{f}^* : \mathbf{H}_n(E) \rightarrow \mathbf{H}_{n+1}(E)$$

because we have  $\mathfrak{f}^* i^* \phi^*(\mathbf{X}_{n-1}(E)) = i^* \phi^* \phi^*(\mathbf{X}_{n-1}(E)) \subset i^* \phi^* \mathbf{X}_n(E)$ . This then induces a  $\phi$ -linear endomorphism  $\mathfrak{f}^* : \mathbf{H}(E) \rightarrow \mathbf{H}(E)$ . Finally, let  $\mathbf{I}_n(E)$  denote the image of  $\partial : \text{Hom}(N^n, \hat{\mathbb{G}}_a) \rightarrow \text{Ext}_A(E, \hat{\mathbb{G}}_a)$ . So  $\text{Hom}(N^n, \hat{\mathbb{G}}_a)/\mathbf{X}_n(E) \simeq \mathbf{I}_n(E)$ . Then  $u$  induces maps  $u^* : \mathbf{I}_n(E) \rightarrow \mathbf{I}_{n+1}(E)$ , and we put  $\mathbf{I}(E) = \lim_{\rightarrow} \mathbf{I}_n(E)$ .

**Proposition 9.3.** *The morphism*

$$u^* : \mathbf{H}_n(E) \otimes K \rightarrow \mathbf{H}_{n+1}(E) \otimes K$$

*is injective. For  $n \geq m$ , it is an isomorphism.*

*Proof.* Consider the following diagram of exact sequences:

$$\begin{array}{ccccccc}
& & 0 & & 0 & & \\
& & \uparrow & & \uparrow & & \\
& & K\langle\phi^{\circ(n-m)^*}\Theta\rangle^\phi & \xrightarrow{i^*\phi^*} & K\langle\Psi_{n+1}\rangle & & \\
& & \uparrow & & \uparrow & & \\
0 & \longrightarrow & \mathbf{X}_n(E)_K^\phi & \xrightarrow{i^*\phi^*} & \mathrm{Hom}_A(N^{n+1}, \hat{\mathbb{G}}_a)_K & \longrightarrow & \mathbf{H}_{n+1}(E)_K \longrightarrow 0 \\
& & \uparrow u^* & & \uparrow u^* & & \uparrow u^* \\
0 & \longrightarrow & \mathbf{X}_{n-1}(E)_K^\phi & \xrightarrow{i^*\phi^*} & \mathrm{Hom}_A(N^n, \hat{\mathbb{G}}_a)_K & \longrightarrow & \mathbf{H}_n(E)_K \longrightarrow 0 \\
& & \uparrow & & \uparrow & & \\
& & 0 & & 0 & & 
\end{array}$$

The cokernels of the two maps  $u^*$  are of the displayed form by propositions 7.4 and 8.6. If  $n < m$ , the expression  $K\langle\phi^{\circ(n-m)^*}\Theta\rangle$  is understood to be zero. The map  $i^*\phi^* : K\langle\phi^{\circ(n-m)^*}\Theta\rangle^\phi \rightarrow K\langle\Psi_{n+1}\rangle$  is injective, by proposition 8.5. Therefore the map  $u^* : \mathbf{H}_n(E)_K \rightarrow \mathbf{H}_{n+1}(E)_K$  is also injective. It is an isomorphism if  $n \geq m$ , because  $K\langle\phi^{\circ(n-m)^*}\Theta\rangle$  is 1-dimensional and hence the map

$$i^*\phi^* : K\langle\phi^{\circ(n-m)^*}\Theta\rangle^\phi \rightarrow K\langle\Psi_{n+1}\rangle$$

is an isomorphism.  $\square$

**Corollary 9.4.** *We have*

$$\mathbf{H}_n(E) \otimes K \simeq \begin{cases} K\langle\Psi_1, \dots, \Psi_n\rangle, & \text{if } n \leq m \\ K\langle\Psi_1, \dots, \Psi_m\rangle, & \text{if } n \geq m \end{cases}$$

Do note that we will promote this to an integral result in (9.7). But before we get there, we will need some preparation.

**Proposition 9.5.** *We have*

$$\mathbf{I}_n(E) \otimes K \simeq \begin{cases} K\langle\Psi_1, \dots, \Psi_n\rangle, & \text{if } n \leq m-1 \\ K\langle\Psi_1, \dots, \Psi_{m-1}\rangle, & \text{if } n \geq m-1 \end{cases}$$

*Proof.* The case  $n \leq m-1$  is clear. So suppose  $n \geq m-1$ . Then  $\mathrm{Hom}_A(N^j, \hat{\mathbb{G}}_a) \otimes K$  has basis  $\Psi_1, \dots, \Psi_j$ , and  $\mathbf{X}_n(E) \otimes K$  has basis  $\Theta_m, \dots, (\phi^{n-m})^*\Theta_m$ . Since each  $(\phi^j)^*\Theta_m$  equals  $\Psi_{m+j}$  plus lower order terms,  $K\langle\Psi_1, \dots, \Psi_{m-1}\rangle$  is a complement to the subspace  $X_n(E)$  of  $\mathrm{Hom}_A(N^n, \hat{\mathbb{G}}_a)$ . Therefore the map  $\partial$  from  $K\langle\Psi_1, \dots, \Psi_{m-1}\rangle$  to the quotient  $I_n(E)$  is an isomorphism.  $\square$

Finally the morphism  $\text{Hom}_A(N^n, \hat{\mathbb{G}}_a) \rightarrow \text{Ext}^\sharp(E, \hat{\mathbb{G}}_a)$  of diagram (9.3) vanishes on  $\phi^*(\mathbf{X}_{n-1}(E))$ , by proposition 9.2, and hence induces a morphism of exact sequences

$$(9.4) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \frac{\mathbf{X}_n(E)}{\phi^*(\mathbf{X}_{n-1}(E)^\phi)} & \longrightarrow & \mathbf{H}_n(E) & \longrightarrow & \mathbf{I}_n(E) \longrightarrow 0 \\ & & \Upsilon \downarrow & & \downarrow \Phi & & \downarrow \\ 0 & \longrightarrow & \text{Lie}(E)^* & \longrightarrow & \text{Ext}^\sharp(E, \hat{\mathbb{G}}_a) & \longrightarrow & \text{Ext}(E, \hat{\mathbb{G}}_a) \longrightarrow 0 \end{array}$$

where as in the introduction,  $\mathbf{I}_n(E)$  denotes the image of  $\partial : \text{Hom}(N^n, \hat{\mathbb{G}}_a) \rightarrow \text{Ext}_A(E, \hat{\mathbb{G}}_a)$ .

**Proposition 9.6.** *The map  $\Phi : \mathbf{H}_n(E) \otimes K \rightarrow \text{Ext}^\sharp(E, \hat{\mathbb{G}}_a) \otimes K$  is injective if and only if  $\gamma \neq 0$ .*

*Proof.* It is enough to show that  $\Upsilon$  is injective if and only if  $\gamma \neq 0$ . By proposition 8.6, the class of  $\Theta_m$  is a  $K$ -linear basis for  $\frac{\mathbf{X}_n(E)}{\phi^*(\mathbf{X}_{n-1}(E)^\phi)} \otimes K$ , and so it is enough to show  $\Phi$  is injective if and only if  $\Upsilon(\Theta_m) \neq 0$ . As in (8.4), write  $\Theta_m = \Psi + g$ . Then by proposition 9.1, it is enough to show  $g'(0) \neq 0$  if and only if  $\gamma \neq 0$ . But this holds because by proposition 8.4, we have  $\gamma = \pi g'(0)$ .  $\square$

**Lemma 9.7.** *Consider the  $\phi$ -linear endomorphism  $F$  of  $K^m$  with matrix*

$$\begin{pmatrix} 0 & 0 & \dots & 0 & \mu_m \\ 1 & 0 & & 0 & \mu_{m-1} \\ 0 & 1 & & 0 & \mu_{m-2} \\ \vdots & & \ddots & \vdots & \vdots \\ 0 & 0 & & 1 & \mu_1 \end{pmatrix},$$

for some given  $\mu_1, \dots, \mu_m \in K$ . If  $K^m$  admits an  $R$ -lattice which is stable under  $F$ , then we have  $\mu_1, \dots, \mu_m \in R$ .

*Proof.* We use Dieudonné–Manin theory. Without loss of generality, we may assume that  $R/\pi R$  is algebraically closed. Let  $P$  denote the polynomial  $F^m - \mu_1 F^{m-1} - \dots - \mu_m$  in the twisted polynomial ring  $K\{F\}$ . Then by (B.1.5) of [Lau] (page 257), there exists an integer  $r \geq 1$  and elements  $\beta_1, \dots, \beta_m \in K(\pi^{1/r})$  such that we have

$$P = (F - \beta_1) \cdots (F - \beta_m)$$

in the ring  $K(\pi^{1/r})\{F\}$  with commutation law  $F\pi^{1/r} = \pi^{1/r}F$ . (Note that the results of [Lau] are stated under the assumption that the residue field of  $R$  is an algebraic closure of  $\mathbb{F}_p$ , but they hold if it is any algebraically closed field of characteristic  $p$ .) Since  $R = K \cap R[\pi^{1/r}]$ , it is enough to show  $\mu_i \in R[\pi^{1/r}]$ . Therefore, by replacing  $R[\pi^{1/r}]$  with  $R$ , it is enough to assume that  $P$  factors as above where in addition all  $\beta_i$  lie in  $K$ .

Now fix  $i$ , and let us show  $\beta_i \in R$ . Assume  $\beta_i \neq 0$ , the case  $\beta_i = 0$  being immediate. Because the (left)  $K\{F\}$ -module  $K^m$  has an  $F$ -stable integral lattice  $M$ , every quotient of  $K^m$  also has a  $F$ -stable integral lattice, namely the image of  $M$ . By (B.1.9) of [Lau] (page 260), for each  $i$ , the  $K\{F\}$ -module  $K^m$  has a quotient (in fact, a summand) isomorphic to  $N = K\{F\}/K\{F\}(F - \pi^{v(\beta_i)})$ . Therefore  $N$

also has a  $F$ -stable integral lattice. But this can happen only if  $v(\beta_i) \geq 0$ , because  $F$  sends the basis element  $1 \in N$  to  $\pi^{v(\beta_i)} \in N$ .  $\square$

**Theorem 9.8.** *If  $E$  splits at  $m$ , then we have  $\lambda_1, \dots, \lambda_{m-1} \in R$ .*

*Proof.* We will prove the cases when  $\gamma = 0$  and  $\gamma \neq 0$  separately.

Case  $\gamma = 0$  When  $\gamma = 0$  we have  $\mathbf{f}^* i^* = i^* \phi^*$ , and hence for all  $n \geq 1$ , this induces a  $\phi$ -linear map  $\mathbf{f}^* : \mathbf{I}_{n-1}(E) \rightarrow \mathbf{I}_n(E)$  as follows

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathbf{X}_n(E) & \xrightarrow{i^*} & \mathrm{Hom}_A(N^n, \hat{\mathbb{G}}_a) & \xrightarrow{\partial} & \mathbf{I}_n(E) \longrightarrow 0 \\ & & \uparrow \phi & & \uparrow \mathbf{f}^* & & \uparrow \mathbf{f}^* \\ 0 & \longrightarrow & \mathbf{X}_{n-1}(E) & \xrightarrow{i^*} & \mathrm{Hom}_A(N^{n-1}, \hat{\mathbb{G}}_a) & \xrightarrow{\partial} & \mathbf{I}_{n-1}(E) \longrightarrow 0 \end{array}$$

Let  $\mathbf{I}(E) = \lim_{\rightarrow} \mathbf{I}_n(E) \subseteq \mathrm{Ext}(E, \hat{\mathbb{G}}_a)$ . Then by proposition 9.5, the vector space  $\mathbf{I}(E)_K$  has a  $K$ -basis  $\partial\Psi_1, \dots, \partial\Psi_{m-1}$ , and with respect to this basis, the  $\phi$ -linear endomorphism  $\mathbf{f}^*$  has matrix

$$\Gamma_0 = \begin{pmatrix} 0 & 0 & \dots & 0 & \lambda_1 \\ 1 & 0 & & 0 & \lambda_2 \\ 0 & 1 & & 0 & \lambda_3 \\ \vdots & & \ddots & \vdots & \vdots \\ 0 & 0 & & 1 & \lambda_{m-1} \end{pmatrix}$$

Since  $\mathbf{I}(E)$  is contained in  $\mathrm{Ext}(E, \hat{\mathbb{G}}_a)$ , it is a finitely generated free  $R$ -module and hence an integral lattice in  $\mathbf{I}(E)_K$ . But then lemma 9.7 implies  $\lambda_1, \dots, \lambda_{m-1} \in R$ .

Case  $\gamma \neq 0$  Let  $\mathbf{H}(E) = \lim_{\rightarrow} \mathbf{H}_n(E)$ . Let us consider the matrix  $\Gamma$  of the  $\phi$ -linear endomorphism  $\mathbf{f}$  of  $\mathbf{H}(E)_K$  with respect to the  $K$ -basis  $\Psi_1, \dots, \Psi_m$  given by corollary 9.4. Let  $\gamma \in R$  be as in proposition 8.4. Then we have

$$\begin{aligned} i^* \phi^* \Theta_m &= \mathbf{f}^* i^* \Theta_m + \gamma \Psi_1 \\ &= \mathbf{f}^* (\Psi_m - \lambda_{m-1} \Psi_{m-1} - \dots - \lambda_1 \Psi_1) + \gamma \Psi_1 \\ &= \mathbf{f}^* (\Psi_m) - \phi(\lambda_{m-1}) \Psi_m - \dots - \phi(\lambda_1) \Psi_2 + \gamma \Psi_1. \end{aligned}$$

Therefore we have

$$\mathbf{f}^* (\Psi_m) \equiv \phi(\lambda_{m-1}) \Psi_m + \dots + \phi(\lambda_1) \Psi_2 - \gamma \Psi_1 \pmod{i^* \phi^* (X_m^\phi)}$$

and hence

$$\Gamma = \begin{pmatrix} 0 & 0 & \dots & 0 & -\gamma \\ 1 & 0 & & 0 & \phi(\lambda_1) \\ 0 & 1 & & 0 & \phi(\lambda_2) \\ \vdots & & \ddots & \vdots & \vdots \\ 0 & 0 & & 0 & \phi(\lambda_{m-2}) \\ 0 & 0 & & 1 & \phi(\lambda_{m-1}) \end{pmatrix}$$

We will now apply lemma 9.7 to the operator  $\mathbf{f}^*$  on  $\mathbf{H}(E)_K$ , but to do this we need to produce an integral lattice  $M$ . Consider the commutative square

$$\begin{array}{ccc} \mathbf{H}(E) & \xrightarrow{\Phi} & \mathrm{Ext}^\sharp(E, \hat{\mathbb{G}}_a) \\ \downarrow & & \downarrow j \\ \mathbf{H}(E)_K & \xrightarrow{\Phi_K} & \mathrm{Ext}^\sharp(E, \hat{\mathbb{G}}_a)_K. \end{array}$$

Let  $M$  denote the image of  $\mathbf{H}(E)$  in  $\mathbf{H}(E)_K$ . It is clearly stable under  $\mathbf{f}^*$ . But also the maps  $\Phi_K$  and  $j$  are injective, by proposition 9.6 and because  $\mathrm{Ext}^\sharp(E, \hat{\mathbb{G}}_a) \simeq R^r$ ; so  $M$  agrees with the image of  $\mathbf{H}(E)$  in  $\mathrm{Ext}^\sharp(E, \hat{\mathbb{G}}_a)$  and is therefore finitely generated.

We can then apply lemma 9.7 and deduce  $\phi(\lambda_{m-1}), \dots, \phi(\lambda_1) \in R$ . This implies  $\lambda_{m-1}, \dots, \lambda_1 \in R$ , since  $R/\pi R$  is a field and hence the Frobenius map on it is injective.  $\square$

**Corollary 9.9.** (1) *The element  $\Theta_m \in \mathbf{X}_m(E)_K$  lies in  $\mathbf{X}_m(E)$ .*  
(2) *For  $n \geq m$ , all the maps in the diagram*

$$\begin{array}{ccc} \mathbf{X}_n(E)/\mathbf{X}_{n-1}(E) & \xrightarrow{\phi^*} & \mathbf{X}_{n+1}(E)/\mathbf{X}_n(E) \\ \downarrow i^* & & \downarrow i^* \\ \mathrm{Hom}_A(N^n, \hat{\mathbb{G}}_a)/\mathrm{Hom}_A(N^{n-1}, \hat{\mathbb{G}}_a) & \xrightarrow{\mathbf{f}^*} & \mathrm{Hom}_A(N^{n+1}, \hat{\mathbb{G}}_a)/\mathrm{Hom}_A(N^n, \hat{\mathbb{G}}_a) \end{array}$$

*are isomorphisms.*

*Proof.* (1): By theorem 9.8, the element  $i^*\Theta_m$  of  $\mathrm{Hom}_A(N^m, \hat{\mathbb{G}}_a)_K$  actually lies in  $\mathrm{Hom}_A(N^m, \hat{\mathbb{G}}_a)$ , and therefore by the exact sequence (8.3) we have  $\Theta_m \in \mathbf{X}_m(E)$ .

(2): By proposition 8.5, we know  $\mathbf{f}^*$  is an isomorphism.

By proposition 8.5, the maps  $i^*$  are injective for all  $n \geq m$ . So to show they are isomorphisms, it is enough to show they are surjective. The  $R$ -linear generator  $\Psi_m$  of  $\mathrm{Hom}_A(N^n, \hat{\mathbb{G}}_a)/\mathrm{Hom}_A(N^{n-1}, \hat{\mathbb{G}}_a)$  is the image of  $\Theta_m$ , which by part (1), lies in  $\mathbf{X}_m(E)$ . Therefore  $i^*$  is surjective for  $n = m$ . Then because  $\mathbf{f}^*$  is an isomorphism, it follows by induction that  $i^*$  is surjective for all  $n \geq m$ .

Finally,  $\phi^*$  is an isomorphism because all the other morphisms in the diagram are.  $\square$

We knew before that  $i^*(\phi^j)^*\Theta_m$  agrees with  $\Psi_{m+j}$  plus lower order rational characters, but the corollary above implies that these lower order characters are in fact integral.

**Theorem 9.10.** *Let  $E$  be a Drinfeld module that splits at  $m$ .*

(1) *For any  $n \geq m$ , the composition*

$$(9.5) \quad \mathbf{X}_n(E) \longrightarrow \mathrm{Hom}_A(N^n, \hat{\mathbb{G}}_a) \longrightarrow \mathrm{Hom}_A(N^n, \hat{\mathbb{G}}_a)/\mathrm{Hom}_A(N^{m-1}, \hat{\mathbb{G}}_a)$$

*is an isomorphism of  $R$ -modules.*

(2)  *$\mathbf{X}_n(E)$  is freely generated as an  $R$ -module by  $\Theta_m, \dots, (\phi^*)^{n-m}\Theta_m$ .*

*Proof.* (i): By corollary 9.9, the induced morphism on each graded piece is an isomorphism. It follows that the map in question is also an isomorphism.

(ii): This follows formally from (i) and the fact, which follows from 9.9, that the map (9.5) sends any  $(\phi^*)^j \Theta_m$  to  $\Psi_{m+j}$  plus lower order terms.  $\square$

**9.2.  $\mathbf{H}(E)$  and de Rham cohomology.** Collecting the results above, we can now prove theorem 1.2. We have isomorphisms

$$R\langle \Psi_1, \dots, \Psi_{m-1} \rangle = \mathrm{Hom}_A(N^{m-1}, \hat{\mathbb{G}}_a) \xrightarrow{\sim} \mathbf{I}_n(E)$$

$$R\langle \Psi_1, \dots, \Psi_m \rangle = \mathrm{Hom}_A(N^m, \hat{\mathbb{G}}_a) \xrightarrow{\sim} \mathbf{H}_n(E)$$

for  $n \geq m$ , and hence in the limit

$$(9.6) \quad R\langle \Psi_1, \dots, \Psi_{m-1} \rangle \xrightarrow{\sim} \mathbf{I}(E)$$

$$(9.7) \quad R\langle \Psi_1, \dots, \Psi_m \rangle \xrightarrow{\sim} \mathbf{H}(E)$$

And so the  $K$ -linear bases of  $K \otimes \mathbf{I}(E)$  and  $K \otimes \mathbf{H}(E)$ —the ones respect to which the action of  $\mathfrak{f}^*$  is described by the matrices  $\Gamma_0$  and  $\Gamma$  in the proof of theorem 9.8—are in fact  $R$ -linear bases of  $\mathbf{I}(E)$  and  $\mathbf{H}(E)$ .

We also have isomorphisms for  $n \geq m$

$$R\langle \Theta_m \rangle = \mathbf{X}_m(E) \xrightarrow{\sim} \mathbf{X}_n(E) / \phi^*(\mathbf{X}_{n-1}(E)^\phi).$$

Combining these, we have the following map between exact sequences of  $R$ -modules, as in (9.4):

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathbf{X}_m(E) & \longrightarrow & \mathbf{H}(E) & \longrightarrow & \mathbf{I}(E) \longrightarrow 0 \\ & & \Upsilon \downarrow & & \downarrow \Phi & & \downarrow \\ 0 & \longrightarrow & \mathrm{Lie}(E)^* & \longrightarrow & \mathrm{Ext}^\sharp(E, \hat{\mathbb{G}}_a) & \longrightarrow & \mathrm{Ext}(E, \hat{\mathbb{G}}_a) \longrightarrow 0 \end{array}$$

where  $\Upsilon$  sends  $\Theta_m$  to  $\gamma/\pi$  (in coordinates). It follows that  $\Phi$  is injective if and only if  $\gamma \neq 0$ .

## 10. COMPUTATION OF $\lambda_1$ AND $\gamma$ IN THE RANK 2 CASE

**Theorem 10.1.** *Let  $A = \mathbb{F}_q[v]$  with  $q \geq 3$ , let  $t \in A$  be an irreducible polynomial of degree  $f$ , and let  $E$  be a Drinfeld module over  $R$  of the form*

$$(10.1) \quad \varphi_E(t)(x) = \pi x + a_1 x^q + a_2 x^{q^2}.$$

Then we have

$$\lambda_1 \equiv (-1)^f w^{\frac{q^f - 1(q^f - 1)}{q - 1}} (1 - a'_1 w^{q^{f-1}} + a'_2 w^{q^{f-1} + q^f})^{q^f - 1} \pmod{\pi},$$

where  $w = a_1 a_2^{-1}$ , and

$$\gamma \pmod{\pi^2} \equiv \begin{cases} \pi \lambda_1 / a_1, & \text{if } \hat{q} \neq q \\ 0, & \text{if } f \geq 3 \text{ or } a_1 \not\equiv 0 \pmod{\pi} \\ -\pi \lambda_1 / a_2, & \text{if } a_1 \equiv 0 \pmod{\pi} \text{ and } f = 2 \end{cases}$$

Observe that when  $\varphi_E(t)(x)$  is of the form  $\pi x + ax^q + x^{q^2}$ , which is always true after changing the coordinate  $x$  (perhaps passing to a cover of  $S$ ), we have the simplified forms

$$(10.2) \quad \lambda_1 \equiv (-1)^f a^{\frac{q^f-1(q^f-1)}{q-1}} (1 - a' a^{q^f-1})^{q^f-1} \pmod{\pi},$$

$$(10.3) \quad \gamma = \pi \lambda_1 / a \pmod{\pi^2}.$$

*Proof.* Let  $\vartheta_1 : N^1 \rightarrow \hat{\mathbb{G}}_a$  be the isomorphism defined in theorem 5.6. Then  $\vartheta_1 \equiv \tau^0 \pmod{\pi}$ . Also  $\vartheta_1$  induces the isomorphism  $(\vartheta_1)_* : \text{Ext}(E, N^1) \rightarrow \text{Ext}(E, \hat{\mathbb{G}}_a)$ . In order to determine the action of  $A$  on  $J^1 E$  and  $J^2 E$  we need to determine how  $t$  acts on the coordinates  $x'$  and  $x''$ . Now we note that  $J^n E \simeq W_n$  can be endowed with the  $\delta$ -coordinates (denoted  $[z, z', z'', \dots]$ ) or the Witt coordinates (denoted  $(z_0, z_1, z_2, \dots)$ ) and they are related by the following in  $J^2 E$  by proposition 3.2

$$(10.4) \quad [z, z', z''] = (z, z', z'' + \pi^{\hat{q}-2}(z')^{\hat{q}})$$

Taking  $\pi$ -derivatives of both sides of equation (10.1) using the formula

$$\delta(ax^{q^j}) = a' x^{\hat{q}q^j} + \phi(a) \pi^{q^j-1} (x')^{q^j},$$

we obtain

$$(10.5) \quad \begin{aligned} \varphi(t)(x') &= \pi' x^{\hat{q}} + a'_1 x^{q\hat{q}} + a'_2 x^{q^2\hat{q}} \\ &\quad + \pi x' + \phi(a_1) \pi^{q-1} (x')^q + \phi(a_2) \pi^{q^2-1} (x')^{q^2} \end{aligned}$$

and

$$(10.6) \quad \begin{aligned} \varphi(t)(x'') &= \pi'' x^{\hat{q}^2} + a''_1 x^{q\hat{q}^2} + a''_2 x^{q^2\hat{q}^2} \\ &\quad + \{\text{terms with } x' \text{ and } x''\} \end{aligned}$$

Then the  $A$ -action  $\varphi_{J^1 E} : A \rightarrow \text{End}(J^1 E)$  is given in Witt coordinates by the  $2 \times 2$  matrix

$$\varphi_{J^1 E}(t) = \begin{pmatrix} \varphi_E(t) & 0 \\ \eta_{J^1 E} & \varphi_{N^1}(t) \end{pmatrix}$$

where  $\eta_{J^1 E} = \pi' x^{\hat{q}} + a'_1 x^{q\hat{q}} + a'_2 x^{q^2\hat{q}}$ . And by (10.6) and (10.4), the  $A$ -action  $A \rightarrow \text{End}(J^2 E)$  is given by the  $(1+2) \times (1+2)$  block matrix

$$\varphi_{J^2 E}(t) = \begin{pmatrix} \varphi_E(t) & 0 \\ \eta_{J^2 E} & \varphi_{N^2}(t) \end{pmatrix}$$

where (using 10.4)  $\eta_{J^2 E}$  is the column vector

$$\eta_{J^2 E} = \begin{pmatrix} \pi' x^{\hat{q}} + a'_1 x^{q\hat{q}} + a'_2 x^{q^2\hat{q}} \\ \Delta(\pi) x^{\hat{q}^2} + \Delta(a_1) x^{q\hat{q}^2} + \Delta(a_2) x^{q^2\hat{q}^2} \end{pmatrix}$$

and where  $\Delta(y) = y'' + \pi^{\hat{q}-2}(y')^{\hat{q}}$ .

Now we will consider two cases—

(1): Consider  $\eta_{\Psi_{1*}(J^1 E)} \in \text{Ext}(E, \hat{\mathbb{G}}_a)$  which is the image of  $\Psi_1$  under the connecting morphism  $\text{Hom}_A(\hat{\mathbb{G}}_a, \hat{\mathbb{G}}_a) \xrightarrow{\partial} \text{Ext}(E, \hat{\mathbb{G}}_a)$  and  $\Psi_1 : N^1 \rightarrow \hat{\mathbb{G}}_a$  is the isomorphism defined in theorem 5.6 and satisfies  $\Psi_1 = \vartheta_1 \circ \mathfrak{f}^0 = \tau^0 \pmod{\pi}$  where



$f^{\circ 0} = \mathbb{1}$ .

$$\begin{array}{ccccccc} 0 & \longrightarrow & N^1 & \longrightarrow & J^1 E & \longrightarrow & E \longrightarrow 0 \\ & & \Psi_1 \downarrow & & \downarrow & & \parallel \\ 0 & \longrightarrow & \hat{\mathbb{G}}_a & \longrightarrow & f_*(J^1 E) & \longrightarrow & E \longrightarrow 0 \end{array}$$

where  $\eta_{J^1 E} = [\pi' x^{\hat{q}} + a'_1 x^{q\hat{q}} + a'_2 x^{q^2 \hat{q}}] \in \text{Ext}(E, N^1)$  Hence

$$\begin{aligned} \eta_{\Psi_{1*}(J^1 E)} &= [\pi' x^{\hat{q}} + a'_1 x^{q\hat{q}} + a'_2 x^{q^2 \hat{q}}] \in \text{Ext}(E, \hat{\mathbb{G}}_a) \\ \partial(\Psi_1) &\equiv [x^{\hat{q}} + a'_1 x^{q\hat{q}} + a'_2 x^{q^2 \hat{q}}] \pmod{\pi}. \end{aligned}$$

(2): Now consider  $\eta_{\Psi_{2*}(J^2 E)} \in \text{Ext}(E, \hat{\mathbb{G}}_a)$  obtained as

$$\begin{array}{ccccccc} 0 & \longrightarrow & N^2 & \longrightarrow & J^2 E & \longrightarrow & E \longrightarrow 0 \\ & & \Psi_2 \downarrow & & \downarrow & & \parallel \\ 0 & \longrightarrow & \hat{\mathbb{G}}_a & \longrightarrow & f_*(J^2 E) & \longrightarrow & E \longrightarrow 0 \end{array}$$

Now we have

$$\eta_{J^2 E} = \left[ \begin{pmatrix} \pi' x^{\hat{q}} + a'_1 x^{q\hat{q}} + a'_2 x^{q^2 \hat{q}} \\ \Delta(\pi)x^{\hat{q}^2} + \Delta(a_1)x^{q\hat{q}^2} + \Delta(a_2)x^{q^2 \hat{q}^2} \end{pmatrix} \right] \in \text{Ext}(E, N^2)$$

Let  $\Xi(y) = (y')^{\hat{q}} + \pi \Delta(y)$ . Then applying  $\Psi_2 = \vartheta_1 \circ f$  and  $f(z_1, z_2) = z_1^{\hat{q}} + \pi z_2$ , we have

$$\begin{aligned} \partial(\Psi_2) &= \eta_{\Psi_{2*}(J^2 E)} = [\vartheta_1(\Xi(\pi)x^{\hat{q}^2} + \Xi(a_1)x^{q\hat{q}^2} + \Xi(a_2)x^{q^2 \hat{q}^2})] \in \text{Ext}(E, \hat{\mathbb{G}}_a) \\ \partial(\Psi_2) &\equiv [\Xi(\pi)x^{\hat{q}^2} + \Xi(a_1)x^{q\hat{q}^2} + \Xi(a_2)x^{q^2 \hat{q}^2}] \pmod{\pi} \\ &\equiv [(\pi')^{\hat{q}}x^{\hat{q}^2} + (a'_1)^{\hat{q}}x^{q\hat{q}^2} + (a'_2)^{\hat{q}}x^{q^2 \hat{q}^2}] \pmod{\pi} \\ &\equiv [x^{\hat{q}^2} + (a'_1)^{\hat{q}}x^{q\hat{q}^2} + (a'_2)^{\hat{q}}x^{q^2 \hat{q}^2}] \pmod{\pi}. \end{aligned}$$

Recall that the map  $R\{\tau\} \rightarrow \text{Ext}(E, \hat{\mathbb{G}}_a)$  given by  $\eta \mapsto [\eta]$  is surjective and the kernel consists of the inner derivations, which is to say all  $\eta$  of the form

$$\pi \alpha - \alpha \circ \varphi_E(t),$$

for some  $\alpha \in R\{\tau\}$ . Let us now work out these relations explicitly for  $\alpha = \tau^0, \tau^1, \tau^2$ . If  $\alpha = \tau^j$ , with  $j \geq 0$ , we get the relation

$$\begin{aligned} \pi \tau^j &= \tau^j (\pi \tau^0 + a_1 \tau^1 + a_2 \tau^2) \\ \tau^{j+2} &= a_2^{-q^j} [(\pi - \pi^{q^j})\tau^j - a_1^{q^j} \tau^{j+1}] \\ \tau^{j+2} &\equiv -(a_1 a_2^{-1})^{q^j} \tau^{j+1} \pmod{\pi} \end{aligned}$$

and hence we have by induction the relations

$$(10.7) \quad \tau^{i+1} \equiv (-1)^i w^{\frac{q^i-1}{q-1}} \tau^1 \pmod{\pi}$$

where  $w = a_1 a_2^{-1}$ , for all  $i \geq 0$ .

Therefore writing  $\hat{q} = q^f$ , we have

$$\begin{aligned}\partial(\Psi_1) &\equiv x^{\hat{q}} + a'_1 x^{q\hat{q}} + a'_2 x^{q^2\hat{q}} \\ &\equiv x^{q^f} + a'_1 x^{q^{f+1}} + a'_2 x^{q^{f+2}} \\ &\equiv \tau^f + a'_1 \tau^{f+1} + a'_2 \tau^{f+2} \\ &\equiv (-1)^{f+1} w^{1+\dots+q^{f-2}} (1 - a'_1 w^{q^{f-1}} + a'_2 w^{q^{f-1}+q^f}) \tau^1\end{aligned}$$

and

$$\begin{aligned}\partial(\Psi_2) &\equiv x^{\hat{q}^2} + (a'_1)^{\hat{q}} x^{q\hat{q}^2} + (a'_2)^{\hat{q}} x^{q^2\hat{q}^2} \\ &\equiv \tau^{2f} + (a'_1)^{q^f} \tau^{2f+1} + (a'_2)^{q^f} \tau^{2f+2} \\ &\equiv (-1)^{2f+1} w^{1+\dots+q^{2f-2}} (1 - (a'_1)^{q^f} w^{q^{2f-1}} + (a'_2)^{q^f} w^{q^{2f-1}+q^{2f}}) \tau^1 \\ &\equiv (-1)^{2f+1} w^{1+\dots+q^{2f-2}} (1 - a'_1 w^{q^{f-1}} + a'_2 w^{q^{f-1}+q^f})^{q^f} \tau^1.\end{aligned}$$

and hence

$$\begin{aligned}\lambda_1 &= \frac{\partial(\Psi_2)}{\partial(\Psi_1)} \equiv (-1)^f w^{q^{f-1}+\dots+q^{2f-2}} (1 - a'_1 w^{q^{f-1}} + a'_2 w^{q^{f-1}+q^f})^{q^f-1} \bmod \pi \\ &\equiv (-1)^f w^{q^{f-1}(1+\dots+q^{f-1})} (1 - a'_1 w^{q^{f-1}} + a'_2 w^{q^{f-1}+q^f})^{q^f-1} \bmod \pi \\ &\equiv (-1)^f w^{\frac{q^{f-1}(q^f-1)}{q-1}} (1 - a'_1 w^{q^{f-1}} + a'_2 w^{q^{f-1}+q^f})^{q^f-1} \bmod \pi\end{aligned}$$

Now we determine  $\gamma$ . Write  $g = \sum_i \alpha_i \tau^i$ . Then from proposition 8.4, we know  $\gamma = \pi \alpha_0$ . Now we will compute  $\alpha_0$ . Let  $(z_0, z_1, z_2) := \varphi_{J^2 E}(t)(x, 0, 0)$ . Then

$$\begin{aligned}\Theta_2(\varphi_{J^2 E}(t)(x, 0, 0)) &= \Psi_2(z_1, z_2) - \lambda_1 \Psi_1(z_1) + g(z_0) \\ &= \vartheta_1(z_1^{\hat{q}} + \pi z_2) - \lambda_1 \vartheta_1(z_1) + g(z_0) \\ &\equiv z_1^{\hat{q}} - \lambda_1 z_1 + g(z_0) \bmod \pi\end{aligned}$$

where  $z_0 = \pi x + a_1 x^q + a_2 x^{q^2}$  and  $z_1 = \pi' x^{\hat{q}} + a'_1 x^{q\hat{q}} + a'_2 x^{q^2\hat{q}}$ . On the other hand from the  $A$ -linearity of  $\Theta_2$  we have

$$\Theta_2(\varphi_{J^2 E}(t)(x, 0, 0)) = \varphi_{\hat{\mathbb{G}}_a}(t) \Theta_2(x, 0, 0) = \pi \Theta_2(x, 0, 0) \equiv 0 \bmod \pi$$

and hence  $z_1^{\hat{q}} - \lambda_1 z_1 + g(z_0) \equiv 0 \bmod \pi$ . Substituting  $z_0$  and  $z_1$  in, we obtain

$$\begin{aligned}0 &\equiv (\pi' x^{\hat{q}} + a'_1 x^{q\hat{q}} + a'_2 x^{q^2\hat{q}})^{\hat{q}} - \lambda_1 (\pi' x^{\hat{q}} + a'_1 x^{q\hat{q}} + a'_2 x^{q^2\hat{q}}) + g(\pi x + a_1 x^q + a_2 x^{q^2}) \\ &\equiv (x^{\hat{q}} + a'_1 x^{q\hat{q}} + a'_2 x^{q^2\hat{q}})^{\hat{q}} - \lambda_1 (x^{\hat{q}} + a'_1 x^{q\hat{q}} + a'_2 x^{q^2\hat{q}}) + g(a_1 x^q + a_2 x^{q^2})\end{aligned}$$

Now substitute  $g(x) = \sum_{j \geq 0} \alpha_j x^{q^j}$  into this and consider the coefficient of  $x^q$ . If  $\hat{q} = q$ , we obtain  $\lambda_1 \equiv \alpha_0 a_1$  and hence

$$\gamma = \pi \alpha_0 \equiv \pi \lambda_1 / a_1 \bmod \pi^2.$$

If  $\hat{q} \neq q$ , we obtain  $\alpha_0 a_1 \equiv 0$  and hence  $\gamma \equiv 0 \bmod \pi^2$  if  $a_1 \not\equiv 0 \bmod \pi$ . If  $a_1 \equiv 0 \bmod \pi$ , we consider the coefficient of  $x^{q^2}$  which is  $\alpha_0 a_2 + \lambda_1$  if  $f = 2$  and  $\alpha_0 a_2$  otherwise. In the case when  $f = 2$  we have  $\alpha_0 \equiv \lambda_1 / a_2 \bmod \pi$  since  $a_2$  is invertible and hence  $\gamma \equiv -\pi \lambda_1 / a_2 \bmod \pi^2$ . When  $f \geq 3$  we have  $\alpha_0 \equiv 0 \bmod \pi$  and hence the result follows.  $\square$

## REFERENCES

- [Bo1] J. Borger: The basic geometry of Witt vectors, I: The affine case, *Algebra and Number Theory* 5 (2011), no. 2, pp 231-285.
- [Bo2] J. Borger: The basic geometry of Witt vectors, II: Spaces, *Mathematische Annalen* 351 (2011), no. 4, pp 877-933.
- [Bui1] A. Buium: Intersections in jet spaces and a conjecture of S.Lang, *Annals of Math.* 136 (1992), 583-593.
- [Bui2] A. Buium: Differential characters of abelian varieties over  $p$ -adic fields, *Invent. Math.*, 122, 2, (1995), 309-340.
- [Bui3] A. Buium: Differential Modular Forms, *Crelle J.*, 520 (2000), 95-167.
- [BP1] A. Buium, B. Poonen: Independence of points on elliptic curves arising from special points on modular and Shimura curves, I: global results, *Duke Math. J.*, 147, 1 (2009), 181-191.
- [BP2] A. Buium, B. Poonen: Independence of points on elliptic curves arising from special points on modular and Shimura curves, II: local results, *Compositio Math.*, 145 (2009), 566-602
- [D76] V.G. Drinfeld, Coverings of  $p$ -adic symmetric domains, *Funkcional. Anal. i Prilozen*, 10(2):29-40, 1976.
- [Ge1] E. -U. Gekeler: de Rham cohomology for Drinfeld modules. *Seminaire de Theorie des Nombres*, Paris 1988/1989, 5785, *Progr. Math.* 91, Birkhauser Boston, Basel, Berlin, 1990.
- [Ge2] E. -U. Gekeler: On the de Rham isomorphism for Drinfeld modules. *J. Reine Angew. Math.* 401 (1989), 188-208.
- [H05] L. Hesselholt, Lecture notes on Witt vectors, *Survey article, unpublished*, 2005.
- [Jo] A. Joyal:  $\delta$ -anneaux et vecteurs de Witt, *C.R. Math. Rep. Acad. Sci. Canada* 7 (1985) 177-182.
- [Lau] G. Laumon: *Cohomology of Drinfeld modular varieties. Part I.* Cambridge University Press, Cambridge (1996)
- [M] J.I. Manin, Rational points of algebraic curves over function fields, *Izvestija Akad. Nauk SSR* 27 (1963), 1395-1440 (in Russian).

*E-mail address:* james.borger@anu.edu.au, arnab.saha@anu.edu.au

AUSTRALIAN NATIONAL UNIVERSITY