

# UNRESTRICTED ALGORITHMS FOR ELEMENTARY AND SPECIAL FUNCTIONS

Invited Paper <sup>1</sup>

**Richard P Brent**

Department of Computer Science  
Australian National University  
Canberra, Australia

We describe some “unrestricted” algorithms which are useful for the computation of elementary and special functions when the precision required is not known in advance. Several general classes of algorithms are identified and illustrated by examples. Applications of such algorithms are mentioned.

## 1 Introduction

Floating-point computations are usually performed with fixed precision: the machine used may have “single” or “double” precision floating-point hardware, or on small machines fixed-precision floating-point operations may be implemented by software or firmware. Most high-level languages support only a small number of floating-point precisions, and those which support an arbitrary number usually demand that the precision be determinable at compile time.

We say that an algorithm has precision  $n$  if its result is computed with error  $O(2^{-n})$ . Usually we are interested in the relative error, but in some cases (e.g. the computation of  $\sin(x)$  for  $x \simeq \pi$ ) it is more appropriate to consider the absolute error.

In certain applications it is desirable that the precision of floating-point operations should be able to be varied at runtime. In this paper we consider algorithms which may be used to evaluate elementary and special functions to precision  $n$ , where  $n$  may be arbitrarily large. Such algorithms have been termed “unrestricted” by Clenshaw and Olver [13]. Note that algorithms which are “unrestricted” in our sense may have domain restrictions (e.g. an “unrestricted” algorithm for  $\exp(x)$  might be applicable only for  $x \geq 0$ ), although such restrictions can often be circumvented by the methods of Section 4, or by combining several algorithms with different domain restrictions.

Unrestricted algorithms depend on the availability of variable-precision floating-point arithmetic. At present this is usually implemented by software, e.g. in the MP package [7], but it could be implemented in firmware or hardware. (Note the historical example of the IBM 1620.)

In the following sections we ignore the possibility of floating-point underflow or overflow. For “ideal” variable-precision arithmetic the exponent range should tend to infinity with the

---

<sup>1</sup>First appeared in *Information Processing 80* (edited by S. H. Lavington), North-Holland, Amsterdam, 1980, 613–619. Retyped with minor corrections by Frances Page at Oxford University Computing Laboratory, 1999. Copyright © 1980, 1999, R. P. Brent and IFIP. rpb052 typeset using L<sup>A</sup>T<sub>E</sub>X.

precision  $n$ . For most purposes, though, a whole-word exponent, as used in MP [7], is adequate to avoid overflow problems.

Applications of variable-precision floating-point arithmetic include:

1. generation and testing of accurate tables of constants (e.g. coefficients in minimax polynomial or rational approximations [9, 25]);
2. computation using numerically unstable algorithms [2, 9];
3. interval arithmetic, where the final intervals may be too large if fixed-precision arithmetic is used [20, 28, 37];
4. truly machine-independent floating-point computations;
5. testing of floating-point hardware for correctness and conformity to standards, e.g. those proposed in [14, 29].
6. number-theoretic computations where very high precision may be essential [24, 30].

In this paper we concentrate on unrestricted algorithms rather than their applications. Section 2 summarises some preliminary results. In each of Sections 3 to 9 we illustrate, by one or two simple examples, a useful general method leading to unrestricted algorithms. Some more specialised methods are mentioned in Section 10. The field is vast and we make no attempt to be comprehensive. For simplicity we usually restrict our attention to real variables and omit details of the rounding error analysis. We also omit any discussion of desirable high-level language facilities to support variable precision arithmetic, for which see [10, 19, 32].

Very few of the algorithms given below are new, in fact most of the identities underlying them may be found in [1] or [35]. What may be new is our viewpoint. Often an excellent unrestricted algorithm is unsuitable for fixed-precision computation, and *vice versa*.

## 2 Basic arithmetic operations

We assume that variable-precision floating-point numbers are represented by an integer exponent and a fraction with  $t$  digits to base  $\beta > 1$ . We call such numbers “precision  $n$ ” numbers if  $n \simeq (t - 1) \log_2 \beta$ . Addition and subtraction of such numbers is straight-forward, and requires  $O(n)$  operations [22, 23]. We assume at least one guard digit, so the relative error in the computed result is at most  $\beta^{1-t} = O(2^{-n})$  (see [36]).

Let  $M(n)$  be the number of operations required for multiplication of precision  $n$  numbers. By the Schönhage-Strassen algorithm [22, 33]

$$M(n) = O(n \log n \log \log n) . \tag{1}$$

For the moderate values of  $n$  which usually arise in applications, an efficient implementation of the classical  $O(n^2)$  algorithm may be faster than the Schönhage-Strassen algorithm or other asymptotically fast algorithms.

Let  $D(n)$  be the number of operations required for division of precision  $n$  numbers. Under plausible assumptions it may be shown that  $D(n) = O(M(n))$  (see, for example, [5]). In practice

the “schoolboy” algorithm, which requires  $O(n^2)$  operations, may be the fastest unless  $n$  is rather large.

It is important to distinguish between multiplication of two precision  $n$  numbers and multiplication or division of a precision  $n$  number by a small (single-precision) integer. The latter require only  $O(n)$  operations if implemented in the obvious way.

### 3 Power series

If  $f(x)$  is analytic in a neighbourhood of some point  $c$ , an obvious method to consider for the evaluation of  $f(x)$  is summation of the Taylor series

$$f(x) = \sum_{j=0}^{k-1} (x-c)^j f^{(j)}(c)/j! + R_k(x, c). \quad (2)$$

As a simple but instructive example we consider the evaluation of  $\exp(x)$  for  $|x| \leq 1$ , using

$$\exp(x) = \sum_{j=0}^{k-1} x^j/j! + R_k(x), \quad (3)$$

where  $|R_k(x)| \leq e/k!$

Using Stirling’s approximation for  $k!$ , we see that  $k \geq K(n) \sim n/\log_2 n$  is sufficient to ensure that  $|R_k(x)| = O(2^{-n})$ . Thus the time required is  $O(nM(n)/\ln n)$ .

In practice it is convenient to sum the series in the forward direction ( $j = 0, 1, \dots, k-1$ ). The terms  $T_j = x/j!$  and partial sums

$$S_j = \sum_{i=0}^j T_i$$

may be generated by the recurrence  $T_j = x \times T_{j-1}/j$ ,  $S_j = S_{j-1} + T_j$ , and the summation terminated when  $|T_k| < 2^{-n}$ . Thus, it is not necessary to estimate  $k$  in advance, as it would be if the series were summed by Horner’s rule in the backward direction ( $j = k-1, k-2, \dots, 0$ ).

We now consider the effect of rounding errors, under the assumption that floating-point operations satisfy

$$fl(x \text{ op } y) = (x \text{ op } y)(1 + \delta), \quad (4)$$

where  $|\delta| \leq \varepsilon$  and “op” = “+”, “−”, “×” or “/”. Here  $\varepsilon \leq \beta^{1-t}$  is the “machine-precision” [36]. Let  $\widehat{T}_j$  be the computed value of  $T_j$ , etc. Thus

$$|\widehat{T}_j - T_j| / |T_j| \leq 2j\varepsilon + O(\varepsilon^2) \quad (5)$$

and

$$\begin{aligned} |\widehat{S}_k - S_k| &\leq k\varepsilon + \sum_{j=1}^k 2j\varepsilon|T_j| + O(\varepsilon^2) \\ &\leq (k+2)e\varepsilon + O(\varepsilon^2) = O(n\varepsilon). \end{aligned} \quad (6)$$

Thus, to get  $|\widehat{S}_k - S_k| = O(2^{-n})$  it is sufficient that  $\varepsilon = O(2^{-n}/n)$ , i.e. we need to work with about  $\log_\beta n$  guard digits. This is not a significant overhead if (as we assume) the number of

digits may vary dynamically. The slightly better error bound obtainable for backward summation is thus of no importance.

In practice it is inefficient to keep  $\varepsilon$  fixed. We can profitably reduce the working precision when computing  $T_k$  from  $T_{k-1}$  if  $|T_{k-1}| \ll 1$ , without significantly increasing the error bound.

It is instructive to consider the effect of relaxing our restriction that  $|x| \leq 1$ . First suppose that  $x$  is large and positive. Since  $|T_j| > |T_{j-1}|$  when  $j < |x|$ , it is clear that the number of terms required in the sum (3) is at least of order  $|x|$ . Thus, the method is slow for large  $|x|$  (see Section 4 for faster methods in this case).

If  $|x|$  is large and  $x$  is negative, the situation is even worse. From Stirling's approximation we have

$$\max_{j \geq 0} |T_j| \simeq \frac{\exp |x|}{\sqrt{2\pi|x|}}, \quad (7)$$

but the result is  $\exp(-|x|)$ , so about  $2|x|/\ln \beta$  guard digits are required to compensate for Lehmer's "catastrophic cancellation" [15]. Since  $\exp(x) = 1/\exp(-x)$ , this problem may easily be avoided, but the corresponding problem is not always so easily avoided for other analytic functions.

In the following sections we generally ignore the effect of rounding errors, but the results obtained above are typical. For an example of an extremely detailed error analysis of an unrestricted algorithm, see [13].

To conclude this section we give a less trivial example where power series expansions are useful. To compute the error function

$$\operatorname{erf}(x) = 2\pi^{-1/2} \int_0^x e^{-u^2} du,$$

we may use the series

$$\operatorname{erf}(x) = 2\pi^{-1/2} \sum_{j=0}^{\infty} \frac{(-1)^j x^{2j+1}}{j!(2j+1)} \quad (8)$$

or

$$\operatorname{erf}(x) = 2\pi^{-1/2} \exp(-x^2) \sum_{j=0}^{\infty} \frac{2^j x^{2j+1}}{1 \cdot 3 \cdot 5 \cdots (2j+1)}. \quad (9)$$

The series (9) is preferable to (8) for moderate  $|x|$  because it involves no cancellation. For large  $|x|$  neither series is satisfactory, because  $\Omega(x^2)$  terms are required, and it is preferable to use the asymptotic expansion or continued fraction for  $\operatorname{erfc}(x) = 1 - \operatorname{erf}(x)$ : see Sections 5 and 6.

## 4 Halving identities

In Section 3 we saw that the power series is not suitable for evaluation of  $\exp(x)$  if  $|x|$  is large. To reduce the size of the argument we may use the identity

$$\exp(x) = [\exp(x/2)]^2 \quad (10)$$

as often as necessary. When applied  $k$  times, (10) gives

$$\exp(x) = [\exp(2^{-k}x)]^{2^k}. \quad (11)$$

If  $k = \lfloor cn^{1/2} \rfloor + \log_2 |x|$  for some positive constant  $c$ , and (11) is used in conjunction with the power series algorithm of Section 3, the time required to evaluate  $\exp(x)$  to precision  $n$  for large  $|x|$  is

$$O[(n^{1/2} + \ln |x|)M(n)] ,$$

better than the  $O[(n/\ln n + |x|)M(n)]$  result of Section 3 (the case  $k = 0$ ).

Similar “halving” (or “doubling”) identities, derived by replacing  $x$  by  $ix$  in (10), may be used to evaluate trigonometric and inverse trigonometric functions [5, 7, 13, 32].

Other identities are useful in special applications: see Section 10 for some examples.

## 5 Asymptotic expansions

Rarely does a single method suffice to evaluate a special function over its whole domain. For example, the exponential integral

$$E_1(x) = \int_x^\infty \frac{\exp(-u)}{u} du \quad (12)$$

is defined for all  $x \neq 0$ . (The Cauchy principal value is taken in (12) if  $x < 0$ .) However, the power series

$$E_1(x) + \gamma + \ln x = \sum_{j=1}^{\infty} \frac{x^j (-1)^{j-1}}{j! j} \quad (13)$$

is unsatisfactory as a means of evaluating  $E_1(x)$  for large positive  $x$ , for the reasons discussed in Section 3 in connection with the power series for  $\exp(x)$ . For sufficiently large  $x$  it is preferable to use the asymptotic expansion [12]

$$E_1(x) = \exp(-x) \sum_{j=1}^k \frac{(j-1)! (-1)^{j-1}}{x^j} + R_k(x) , \quad (14)$$

where

$$R_k(x) = k! (-1)^k \int_x^\infty \frac{\exp(-u)}{u^{k+1}} du . \quad (15)$$

For large positive  $x$ , the relative error attainable by using (14) with  $k \simeq x$  is  $O(x^{1/2} \exp(-x))$ , because

$$|R_k(k)| \leq k! k^{-(k+1)} \exp(-k) = O(k^{-1/2} \exp(-2k)) . \quad (16)$$

Thus, the asymptotic series may be used to evaluate  $E_1(x)$  to precision  $n$  when  $x > n \ln 2 + O(\ln n)$ . (Similarly if  $(-x) > n \ln 2 + O(\ln n)$ , although the estimation of  $|R_k(-k)|$  is more difficult than that of  $|R_k(k)|$ .)

There are many other examples where asymptotic expansions are useful, e.g. for  $\operatorname{erfc}(x)$  (mentioned in Section 3), for Bessel functions [11, 35], etc. Asymptotic expansions often arise when the convergence of series is accelerated by the Euler-Maclaurin sum formula [1]. For example, the Riemann zeta function  $\zeta(s)$  is defined for  $R(s) > 1$  by

$$\zeta(s) = \sum_{j=1}^{\infty} j^{-s} , \quad (17)$$

and by analytic continuation for other  $s \neq 1$ . (Here we allow complex  $s$ .)  $\zeta(s)$  may be evaluated to any desired precision if  $m$  and  $p$  are chosen large enough in the Euler-Maclaurin formula [8]

$$\zeta(s) = \sum_{j=1}^{p-1} j^{-s} + \frac{1}{2}p^{-s} + \frac{p^{1-s}}{s-1} + \sum_{k=1}^m T_{k,p}(s) + E_{m,p}(s), \quad (18)$$

where

$$T_{k,p}(s) = \frac{B_{2k}}{(2k)!} p^{1-s-2k} \prod_{j=0}^{2k-2} (s+j), \quad (19)$$

$$|E_{m,p}(s)| < |T_{m+1,p}(s) (s+2m+1)/(\sigma+2m+1)|, \quad (20)$$

$m \geq 0$ ,  $p \geq 1$ ,  $\sigma = R(s) > -(2m+1)$ , and the  $B_{2k}$  are Bernoulli numbers.

In arbitrary-precision computations we must be able to compute as many terms of an asymptotic expansion as are required to give the desired accuracy. It is easy to see that  $m$  in (18) can not be bounded as the precision  $n \rightarrow \infty$ , else  $p$  would have to increase as an exponential function of  $n$ . To evaluate  $\zeta(s)$  from (18) to precision  $n$  in time polynomial in  $n$ , both  $m$  and  $p$  must tend to infinity with  $n$ . Thus, the Bernoulli numbers  $B_2, \dots, B_{2m}$  can not be stored in a table of fixed size, but must be computed when needed (see Sections 7 and 9). For this reason we can not use asymptotic expansions when the general form of the coefficients is unknown (such as Stirling's formula for  $\Gamma(x)$ ) in arbitrary-precision calculations. Often there is a related expansion with known coefficients, e.g. the asymptotic expansion for  $\ln \Gamma(x)$  has coefficients related to the Bernoulli numbers, like (19).

## 6 Continued fractions

Sometimes continued fractions are preferable to power series or asymptotic expansions. For example, Euler's continued fraction [34]

$$\exp(x) E_1(x) = 1/x + 1/1 + 1/x + 2/1 + 2/x + 3/1 + \dots \quad (21)$$

converges for all real  $x > 0$ , and is better for computation of  $E_1(x)$  than the power series (13) in the region where the power series suffers from catastrophic cancellation but the asymptotic expansion (14) is not sufficiently accurate. Convergence of (21) is slow if  $x$  is small, so (21) is preferred for precision  $n$  evaluation of  $E_1(x)$  only when  $x \in (c_1 n, c_2 n)$ ,  $c_1 \simeq 0.1$ ,  $c_2 \simeq \ln 2$ .

It is well known that continued fractions may be evaluated by either forward or backward recurrence relations. Consider the finite continued fraction

$$y = a_1/b_1 + a_2/b_2 + \dots + a_k/b_k. \quad (22)$$

The backward recurrence is  $R_k = 1$ ,  $R_{k-1} = b_k$ ,

$$R_j = b_{j+1} R_{j+1} + a_{j+2} R_{j+2} \quad (j = k-2, \dots, 0), \quad (23)$$

and  $y = a_1 R_1 / R_0$ . The forward recurrence is  $P_0 = 0$ ,  $P_1 = a_1$ ,  $Q_0 = 1$ ,  $Q_1 = b_1$ ,

$$\left. \begin{aligned} P_j &= b_j P_{j-1} + a_j P_{j-2} \\ Q_j &= b_j Q_{j-1} + a_j Q_{j-2} \end{aligned} \right\} \quad (j = 2, \dots, k), \quad (24)$$

and  $y = P_k / Q_k$ .

The advantage of evaluating an infinite continued fraction such as (21) via the forward recurrence is that  $k$  need not be chosen in advance; we can stop when  $|D_k|$  is sufficiently small, where

$$D_k = \frac{P_k}{Q_k} - \frac{P_{k-1}}{Q_{k-1}}. \quad (25)$$

The disadvantage of the forward recurrence is that twice as many arithmetic operations are required as for the backward recurrence with the same value of  $k$ . There is a simple solution to this dilemma if we are working with variable-precision floating-point arithmetic which is much more expensive than single-precision floating-point. We use the forward recurrence with single-precision arithmetic (scaled to avoid overflow/underflow) to estimate  $k$ , then use the backward recurrence with variable-precision arithmetic. One trick is needed: to evaluate  $D_k$  using scaled single-precision we use the recurrence

$$\left. \begin{aligned} D_1 &= a_1/b_1, \\ D_j &= -a_j Q_{j-2} D_{j-1} / Q_j \quad (j = 2, 3, \dots) \end{aligned} \right\} \quad (26)$$

which avoids the cancellation inherent in (25).

In recent versions of the MP package [7] we have used the continued fraction (21) in the manner just described, and similar continued fractions could well be used for the computation of other special functions. Since power series and asymptotic series are generally easier to analyse and program than continued fractions, we have avoided continued fractions except where they are clearly superior to the other methods.

## 7 Recurrence relations

The evaluation of special functions by continued fractions is a special case of their evaluation by recurrence relations. For example, the Bessel functions  $J_\nu(x)$  satisfy the recurrence relation

$$J_{\nu-1}(x) + J_{\nu+1}(x) = \frac{2\nu}{x} J_\nu(x) \quad (27)$$

which may be evaluated backwards (compare (23)), using a normalisation condition such as

$$J_0(x) + 2 \sum_{\nu=1}^{\infty} J_{2\nu}(x) = 1. \quad (28)$$

This seems to be the most effective method in the region where Hankel's asymptotic expansion is insufficiently accurate but the power series

$$J_\nu(x) = \left(\frac{x}{2}\right)^\nu \sum_{j=0}^{\infty} \frac{(-x^2/4)^j}{j! \Gamma(\nu + j + 1)} \quad (29)$$

suffers from catastrophic cancellation. For details see [17].

In Section 5 the constants  $C_k = B_{2k}/(2k)!$  were required, where the  $B_{2k}$  are Bernoulli numbers. The  $C_k$  are defined by the generating function

$$\sum_{k=0}^{\infty} C_k x^{2k} = \frac{x}{e^x - 1} + \frac{x}{2}. \quad (30)$$

Multiplying both sides by  $e^x - 1$  and equating coefficients gives the recurrence relation

$$\frac{C_k}{1!} + \frac{C_{k-1}}{3!} + \cdots + \frac{C_1}{(2k-1)!} = \frac{k - \frac{1}{2}}{(2k+1)!}, \quad (31)$$

which has often been used to evaluate Bernoulli numbers [21].

Unfortunately, forward evaluation of the recurrence (31) is numerically unstable: using precision  $n$  the relative error in the computed  $C_k$  is of order  $4^k 2^{-n}$ . We shall not prove this, but shall indicate why such behaviour is to be expected. Consider the ‘‘homogeneous’’ recurrence

$$\frac{\widehat{C}_k}{1!} + \frac{\widehat{C}_{k-1}}{3!} + \cdots + \frac{\widehat{C}_1}{(2k-1)!} = 0 \quad (k \geq 2) \quad (32)$$

with  $\widehat{C}_1 = 1$ , and let

$$\widehat{G}(x) = \sum_{k=1}^{\infty} \widehat{C}_k x^{2k} \quad (33)$$

be the generating function for the  $\widehat{C}_k$ . It is easy to show that

$$\widehat{G}(x) = \frac{x^3}{\sinh x}. \quad (34)$$

Thus  $\widehat{G}(x)$  has poles at  $\pm i\pi$ , and

$$|\widehat{C}_k| \geq K \pi^{-2k} \quad (35)$$

for some  $K > 0$ . This suggests that an error of order  $2^{-n}$  in an early value of  $C_j$  propagates to give an (absolute) error of order  $2^{-n} \pi^{-2k}$  in  $C_k$  for large  $k$ . Since  $|C_k| \sim (2\pi)^{-2k}$ , this absolute error corresponds to a relative error of order  $2^{2k-n} = 4^k 2^{-n}$  in  $C_k$ .

Despite its numerical instability, use of (31) may give the  $C_k$  to acceptable accuracy if they are only needed to generate coefficients in an Euler-Maclaurin expansion whose successive terms diminish by at least a factor of 4. If the  $C_k$  or  $B_{2k}$  are required to precision  $n$ , either (31) must be used with sufficient guard digits, or a more stable recurrence must be used. If we multiply both sides of (30) by  $\sinh(x/2)/x$  and equate coefficients, we get the recurrence

$$C_k + \frac{C_{k-1}}{3! 4} + \cdots + \frac{C_1}{(2k-1)! 4^{k-1}} = \frac{2k}{(2k+1)! 4^k} \quad (36)$$

If (36) is used to evaluate  $C_k$ , using precision  $n$  arithmetic, the error is only  $O(k^2 2^{-n})$ . Thus, this method is currently used in the MP package instead of a method based on (31).

## 8 Newton’s method

Newton’s method and related zero-finding methods may be used to evaluate a function if we have an algorithm for evaluation of the inverse function. For example, applying Newton’s method to  $f(x) = y - x^{-m}$  (where  $y$  is regarded as constant) gives the iteration

$$x_{j+1} = x_j + x_j(1 - x_j^m y)/m, \quad (37)$$

which converges (from a sufficiently good initial approximation) to  $y^{-1/m}$ . Note that (37) does not involve divisions except by the small integer  $m$ .



Similarly, applying Newton's method to  $f(x) = \exp(x) - y$  gives the iteration

$$x_{j+1} = x_j + y \exp(-x_j) - 1 . \quad (38)$$

which converges to  $\ln y$  if  $x_0$  is a sufficiently good initial approximation.

Newton's method generally has second order convergence, so we may start with low precision and approximately double it at each iteration. Thus, the work required is of the same order as the work for the final iteration. Applied to (37) with  $m = 1$  and 2, this argument shows that reciprocals and square roots can be found to precision  $n$  in  $O(M(n))$  operations. For further details, and a comparison of the efficiencies of various root-finding methods for variable-precision computations, see [4, 5].

## 9 Contour integration

In this section we assume that facilities for variable-precision complex arithmetic are available. Let  $f(z)$  be holomorphic in the disc  $|z| < R$ ,  $R > 1$ , and let the power series for  $f$  be

$$f(z) = \sum_{j=0}^{\infty} a_j z^j \quad (39)$$

From Cauchy's theorem [18] we have

$$a_j = \frac{1}{2\pi i} \int_C \frac{f(z)}{z^{j+1}} dz , \quad (40)$$

where  $C$  is the unit circle. The contour integral in (40) may be approximated numerically by sums

$$S_{j,k} = \frac{1}{k} \sum_{m=0}^{k-1} f(e^{2\pi i m/k}) e^{-2\pi i j m/k} . \quad (41)$$

From Cauchy's theorem, provided  $j < k$  and the contour  $C$  is enlarged slightly to enclose the  $k$ -th roots of unity, we have

$$\begin{aligned} S_{j,k} - a_j &= \frac{1}{2\pi i} \int_C \frac{f(z)}{(z^k - 1)z^{j+1}} dz \\ &= a_{j+k} + a_{j+2k} + \dots , \end{aligned} \quad (42)$$

so  $|S_{j,k} - a_j| = O((R - \delta)^{-(j+k)})$  as  $k \rightarrow \infty$ , for any  $\delta > 0$ .

For example, let

$$f(z) = \frac{z}{e^z - 1} + \frac{z}{2} \quad (43)$$

as in Section 7, so  $a_{2j} = B_{2j}/(2j)!$  and  $R = 2\pi$ . Then

$$S_{2j,k} - \frac{B_{2j}}{(2j)!} = \frac{B_{2j+k}}{(2j+k)!} + \frac{B_{2j+2k}}{(2j+2k)!} + \dots , \quad (44)$$

so we can evaluate  $B_{2j}$  with relative error  $O((2\pi)^{-k})$  by evaluating  $f(z)$  at  $k$  points on the unit circle. (By symmetry and conjugacy only  $k/4 + 1$  evaluations are required if  $k$  is a multiple of four.) If  $\exp(-2\pi i j m/k)$  is computed efficiently from  $\exp(-2\pi i/k)$  in the obvious way, the time required to evaluate  $B_2, \dots, B_{2j}$  to precision  $n$  is  $O(jnM(n))$ , and the space required is  $O(n)$ . The recurrence relation method of Section 7 requires time only  $O(j^2n)$ , but space  $O(jn)$ . Thus, the method of contour integration is recommended if space is more important than time.

For further discussion of the contour integration method, see [26].

## 10 Special methods

In this section we mention two of a large number of “special” methods which are useful but less generally applicable than the methods of Sections 3 to 9. The first such method is the conversion of a power series which suffers from catastrophic cancellation to one which is better behaved numerically. One example, (9), has already been given. Another example occurs with

$$E(x) = \int_0^x \frac{(1 - e^{-u})}{u} du = \sum_{j=1}^{\infty} \frac{x^j (-1)^{j-1}}{j! j} \quad (45)$$

(a series encountered in Section 5). Multiplying by  $\exp(x)$  and using some well known identities, we find

$$\exp(x)E(x) = \sum_{j=1}^{\infty} H_j x^j / j! , \quad (46)$$

where

$$H_j = \sum_{m=1}^j \frac{1}{m} . \quad (47)$$

If  $x$  is large and positive, the series in (46) is much better behaved numerically than the series in (45). For an application where  $E(x)$  was required to high precision with  $x$  a positive integer, see [11]. At first sight it appears that, in this application, the summation to precision  $n$  of  $k$  terms in the series (45) requires  $O(kn)$  operations, while (46) requires  $\Omega(kM(n))$  operations. However, by a “summation by parts” trick described in [11], this can be reduced to  $O(kn)$  operations.

Our second “special” method is the evaluation of  $\pi$  and elementary functions by the arithmetic-geometric mean (AGM) iteration. It is well known that the AGM can be used to compute elliptic integrals, but perhaps less well known that it can also be used to compute  $\pi$  and elementary functions, and gives the fastest known methods when the precision  $n$  is very large [4, 6].

The AGM of two positive numbers  $a_0$  and  $b_0$  is  $a = \lim_{j \rightarrow \infty} a_j = \lim_{j \rightarrow \infty} b_j$ , where

$$a_{j+1} = \frac{a_j + b_j}{2} \quad (48)$$

and

$$b_{j+1} = \sqrt{a_j b_j} . \quad (49)$$

There is no essential loss of generality in assuming that  $a_0 = 1$  and  $b_0 = \cos \phi$ . Gauss [16] showed that  $2a = \pi/K(\phi)$ , where

$$K(\phi) = \int_0^{\pi/2} (1 - \sin^2 \phi \sin^2 \theta)^{-1/2} d\theta \quad (50)$$

is the complete elliptic integral of the first kind. A simple proof is given in [27].

The AGM iteration converges quadratically: if  $\varepsilon_j = 1 - b_j/a_j$  then

$$\varepsilon_{j+1} = 1 - 2(1 - \varepsilon_j)^{1/2}/(2 - \varepsilon_j) = \varepsilon_j^2/8 + O(\varepsilon_j^3) . \quad (51)$$

Using the AGM and an identity of Lagrange, we get a family of quadratically convergent algorithms for the computation of  $\pi$ . The simplest of these is:

```

a := 1 ;
b := 1/√2 ;
t := 1/4 ;
j := 1 ;
repeat
    y := a ;
    a := (a + b)/2 ;
    b := √(b × y) ;
    t := t - j × (a - y)2 ;
    j := 2 × j
until (a - b) < tolerance ;
return a2/t.

```

After  $k$  iterations the error  $|a^2/t - \pi|$  is about  $8\pi \exp(-2^k\pi)$ , e.g.  $k = 5$  gives error less than  $10^{-42}$ . For further details see [6, 31].

In [3, 4, 6] it is shown how the AGM may be used to compute the elementary functions  $\exp(x)$ ,  $\ln(x)$ ,  $\operatorname{atan}(x)$ ,  $\sin(x)$  etc. to precision  $n$  in  $O(M(n) \log n)$  operations. The factor “ $\log n$ ” arises because  $O(\log n)$  iterations of the AGM are required. It is important to note that the AGM iteration is not self-correcting, so the trick of starting with low precision and doubling it on each iteration (as used in Section 8) is not applicable.

## 11 Summary

Many “classical” methods may be adapted for use in variable-precision computations; others are not readily adaptable. Since the performance criteria are different in variable-precision applications, the best method may be one which is not well-suited to fixed-precision computations. For example, it might be numerically unstable, and thus require the working precision to be increased. The examples given in Sections 3 to 10 above are intended to illustrate the main ideas of variable-precision algorithms.

## 12 Acknowledgement

Christian Reinsch kindly suggested that the use of (36) would be faster than the method described in Section 6.11 of [7].

## References

- [1] M.A. Abramowitz and I.A. Stegun (eds.), *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, National Bureau of Standards, Washington, D.C., 1964 (reprinted by Dover, New York, 1965).
- [2] R.E. Bank and D.J. Rose, “Extrapolated fast direct algorithms for elliptic boundary value problems”, in *Algorithms and Complexity* (ed. by J.F. Traub), Academic Press, New York, 1976, 201–249.
- [3] M. Beeler, R.W. Gosper and R. Schroepfel, *Hakmem*, M.I.T. Artificial Intelligence Lab. Memo No. 239, Feb. 1972. [Available from <http://www.inwap.com/pdp10/hbaker/hakmem/hakmem.html> ]
- [4] R.P. Brent, “Multiple-precision zero-finding methods and the complexity of elementary function evaluation”, in *Analytic Computational Complexity* (ed. by J.F. Traub), Academic Press, New York, 1975, 151–176. [Available from <ftp://ftp.comlab.ox.ac.uk/pub/Documents/techpapers/Richard.Brent/rpb028.ps.gz> ]
- [5] R.P. Brent, “The complexity of multiple-precision arithmetic”, in *The Complexity of Computational Problem Solving* (eds. R.S. Anderssen and R.P. Brent), Queensland Univ. Press, Brisbane, 1976, 126–165. [Available from <ftp://ftp.comlab.ox.ac.uk/pub/Documents/techpapers/Richard.Brent/rpb032.ps.gz> ]
- [6] R.P. Brent, “Fast multiple-precision evaluation of elementary functions”, *J.ACM* 23 (1976), 242–251.
- [7] R.P. Brent, “A Fortran multiple-precision arithmetic package”, *ACM Trans. Math. Software* 4 (1978), 57–70.
- [8] R.P. Brent, “On the zeros of the Riemann zeta function in the critical strip”, *Math. Comp.* 33 (1979), 1361–1372.
- [9] R.P. Brent, *Numerical investigation of the Riemann-Siegel approximation*, unpublished notes, 1979.
- [10] R.P. Brent, J.A. Hooper and J.M. Yohe, “An Augment interface for Brent’s multiple-precision arithmetic package”, *ACM Trans. Math. Software*, to appear. [Appeared in vol. 6 (1980), 146–149.]
- [11] R.P. Brent and E.M. McMillan, “Some new algorithms for high-precision computation of Euler’s constant”, *Math. Comp.*, to appear. [Appeared in vol. 34 (1980), 305–312.]
- [12] N.G. de Bruijn, *Asymptotic Methods in Analysis*, 3rd edition, North-Holland, 1970.
- [13] C.W. Clenshaw and F.W.J. Olver, “An unrestricted algorithm for the exponential function”, *SIAM J. Numer. Anal.*, to appear. [Appeared in vol. 17, 1980, 310–331.]
- [14] J.T. Coonen, W. Kahan, J. Palmer, T. Pittman and D. Stevenson, “A proposed standard for binary floating point arithmetic, draft 5.11”, *ACM SIGNUM Newsletter*, October 1979, 4–12.
- [15] G.E. Forsythe, “Pitfalls in computation, or why a math book isn’t enough”, *Amer. Math. Monthly* 77 (1970), 931–956.
- [16] C.F. Gauss, *Carl Friedrich Gauss Werke*, Bd. 3, Göttingen, 1876, 362–403.
- [17] W. Gautschi, “Algorithm 236: Bessel functions of the first kind”, *Comm. ACM* 7 (1964), 479–480.
- [18] E. Hille, *Analytic Function Theory*, Vol. 1, Blaisdell, New York, 1959, Ch. 7.
- [19] T.E. Hull and J.J. Hofbauer, *Language facilities for multiple-precision floating-point computation*, Dept. of Computer Science, Univ. of Toronto, 1974.
- [20] J.P. Jeter, *A Variable-Precision Interval Data Type Extension to Fortran*, M.Sc. thesis, Dept. of Computer Science, Univ. of S.W. Louisiana, Lafayette, Louisiana, July 1979.
- [21] D.E. Knuth, “Euler’s constant to 1271 places”, *Math. Comp.* 16 (1962), 275–281.

- [22] D.E. Knuth, *The Art of Computer Programming*, Vol. 2, Addison Wesley, Reading, Mass., 1969.
- [23] D.E. Knuth, “Big Omicron and Big Omega and Big Theta”, *SIGACT News* 8, 2 (1976), 18–24.
- [24] D.H. Lehmer, “Tables to many places of decimals”, *Math. Tables Aids Comput.* 1 (1943), 30–31. [The journal is now called *Mathematics of Computation*.]
- [25] Y.L. Luke, *Algorithms for the Computation of Mathematical Functions*, Academic Press, New York, 1977.
- [26] J.N. Lyness and C.B. Moler, “Numerical differentiation of analytic functions”, *SIAM J. Numer. Anal.* 4 (1967), 202–210.
- [27] Z.A. Melzak, *Companion to Concrete Mathematics*, Wiley, New York, 1973.
- [28] R.E. Moore, *Interval Analysis*, Prentice-Hall, New Jersey, 1966.
- [29] M. Payne and W. Strecker, “Draft proposal for a binary normalized floating point standard”, *ACM SIGNUM Newsletter*, October 1979, 24–30.
- [30] H.J.J. te Riele, “Computations concerning the conjecture of Mertens”, *J. reine angew. Math.* 311/312 (1979), 356–360.
- [31] E. Salamin, “Computation of  $\pi$  using arithmetic-geometric mean”, *Math. Comp.* 30 (1976), 565–570.
- [32] J.L. Schonfelder and J.T. Thomason, *Applications support by direct language extension – an arbitrary precision arithmetic facility in Algol 68*, Comp. Centre, Univ. of Birmingham, Birmingham, U.K., 1975.
- [33] A. Schönhage and V. Strassen, “Schnelle Multiplikation grosser Zahlen”, *Computing* 7 (1971), 281–292.
- [34] H. Wall, *Analytic Theory of Continued Fractions*, Van Nostrand, New York, 1948.
- [35] E.T. Whittaker and G.N. Watson, *A Course of Modern Analysis*, Cambridge Univ. Press, 1902.
- [36] J.H. Wilkinson, *Rounding Errors in Algebraic Processes*, HMSO, London, 1963.
- [37] J.M. Yohe, *The interval arithmetic package – multiple precision version*, MRC Tech. Summary Report No. 1908, Math. Res. Center, Madison, Jan. 1979.